

DAFTAR PUSTAKA

- Allakhverdi Z., Delespesse G., 2012, *Hematopoietic progenitor cells are innate Th2 cytokine-producing cells*, *Allergy* 67: 4–9.
- Baratawidjaja G. K., Rengganis I., 2014, *Imunologi Dasar*, Edisi ke-11, Badan Penerbit FKUI, Jakarta, 217-85.
- Bongso A & Lee EH. 2005. *Stem Cells: From Bench to Bedside*. Singapore: World Scientific Publishing Co. Pte. Ltd.
- Cosgun KN. *Cell Stem Cell* 15: 227-238, 2014
- Crisostomo PR, Wang Y, Markel TA, Wang M, Lahm T, et al. (2008) Human mesenchymal stem cells stimulated by TNF-alpha, LPS, or hypoxia produce growth factors by an NF kappa B- but not JNK-dependent mechanism. *Am J Physiol Cell Physiol* 294: C675–682.
- Dani Halim, Harry Murti, Ferry Sandra, Areif boediono, Tono Djuwantono, Boenjamin setiawan. *Stem Cell, dasar teori & aplikasi klinis*, 2010.4:109.
- Dressel R, Schindehütte J, Kuhlmann T, Elsner L, Novota P, Baier P, et al. 2008. *The tumorigenicity of mouse embryonic stem cells and in vitro differentiated neuronal cells is controlled by the recipients' immune response*. PLoS ONE.
- Durum and Muege.2003. *Hamster Fed Diet High Saturated Fat have Increased Cholesterol accumulated and cytokine Production in Aortic Area compared with Cholesterol Fed Hamster with Moderately Elevated plasma non HDL Cholesterol Concentration*. *JournalNutrisi Immunology*. Universitas of Massachusetts.
- Fogg D, Sibon C, Miled C, Jung S, Aucouturier P, Littman D, et al. *Aclonogenic bone marrow progenitro spesific for macrophages and dendritic cells*. *Science*. 2006;311:83-7 PMID16322423.
- Granick JL., Simon SI., Borjesson DL., 2012, *Hematopoietic stem and progenitor cells as effectors in innate immunity*, *Bone Marrow Res*: 165107.
- J, Jiang., R, n Goel., MA, Iftekhar., R, Visaria., JD,et al. 2008. *TNF alpha induced accentuation in cryoinjury: mechanism in vitro and in vivo*. USA: US National Library of Medicine National Institutes of Health.
- Kang S.K.I. 2012. *Journey of Mesenchymal Stem Cells for Homing :strategies to enhance efficacy and safety of stem cell therapy*. stem cells international

- Kitagawa M., Saito I., Kuwata T., Yoshida S., Yamaguchi S., Takahashi M., et al., 2007, *Overexpression of tumor necrosis factor (TNF)-alpha and interferon (IFN)-gamma by bone marrow cells from patients with myelodysplastic syndromes*, *Leukemia*;11:2049–2054.
- Kumar V., Abbas AK., Fausto N. 2005. Neoplasia. In: Robbins and Cotran *Pathology Basis of Disease*. 7th Ed, Philadelphia. Elsevier Saunders.:1041- 1042.
- Kwon YW., Heo SC., Jeong GO., Yoon JW., Mo WM., Lee MJ., et al., 2013, *Tumor necrosis factor- α -activated mesenchymal stem cells promote endothelial progenitor cell homing and angiogenesis*, *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 2136–2144.
- Lagasse, E., Connors, H., Al Dhalimy, M., Reitsma, M., Dohse, M., Osborne, L., Wang, X., Finegold, M., Weissman, I.L., and Grompe, M. (2007). *Purified hematopoietic stem cells can differentiate into hepatocytes in vivo*. *Nat. Med.* 6, 1229–1234.
- Lagasse, E., Connors, H., Al Dhalimy, M., Reitsma, M., Dohse, M., Osborne, L., Wang, X., Finegold, M., Weissman, I.L., and Grompe, M., 2007, *Purified hematopoietic stem cells can differentiate into hepatocytes in vivo*, *Nat. Med.* 6, 1229–1234.
- Locksley RM., Killeen N., Lenardo MJ., 2001, *The TNF and TNF receptor superfamilies: integrating mammalian biology*, *Cell*, 104; 487–501.
- Rodriguez, L.G., Wu X., and Guan J.L. 2005. *Wound-healing assay*. *Mol Biol* 294: 23-9.
- Setiawan B. 2006. *Aplikasi terapeutik sel stem embrionik pada berbagai penyakit degeneratif*. *Cermin Dunia Kedokteran*
- Shenghui H, Nakada D, and Morrison SJ. 2009. *Mechanisms of Stem cell Self-Renewal* *Annu. Rev. Cell Dev. Biol.*
- Singhal S, Vachani A, Antin-Ozerkis A, Kaiser LR, Albelda SM. *Prognostic implication of cell cycle, apoptosis, and angiogenesis biomarkers in non-small cell lung cancer: a review*. *Clin Cancer Res* 2005; 11:3974-86
- Sukhanov, S., Higashi, Y., Yung Shai, S., Vaughn, C., Mohler, J., Li, Y., Hua Song, Y., Titterton, J., Delafontaine, P. 2007. *IGF-1 Reduces Inflammatory Responses, suppresses Oxidative Stress, and Decreases Atherosclerosis Progression in ApoE-Deficient Mice*. *Arterioscler Thromb Vasc Biol.* 27;84-2690.
- WHO (World Health Organization), *Global Status Report On Noncommunicable diseases* 2014

- Wilmana, P.F., dan Gan, S., 2007. Analgesik-Antipiretik Analgesik Anti-Inflamasi Nonsteroid dan Obat Gangguan Sendi Lainnya. *Dalam: Gan, S.,Setiabudy, R., dan Elysabeth, eds. Farmakologi dan Terapi*. Edisi 5. Jakarta:Departemen Farmakologi dan Terapeutik FK UI, 237-239.
- Yoshida T., Hakuba N., Morizane I., Fujita K., Cao F., et al., 2007, *Hematopoietic stem cells prevent hair cell death after transient cochlear ischemia through paracrine effects*, *Neuroscience* 145: 923–930.
- Chen L., Tredget EE., Wu PY., Wu Y., 2008, “*Paracrine factors of mesenchymal stem cells recruit macrophages and endothelial lineage cells and enhance wound healing*”, *PLoS One*; 3:e1886.
- Fu X., Han B., Cai S., et al., 2009, “*Migration of bone marrow-derived mesenchymal stem cells induced by tumor necrosis factor-alpha and its possible role in wound healing*”, *Wound Repair Regen*; 17(2):185-191.
- Gutiérrez FM., Rodríguez FB., Alvarez GJ., et al., 2011, “*Functional recovery after hematic administration of allogenic mesenchymal stem cells in acute ischemic stroke in rats*”, *Neuroscience*; 175: 394-405.
- Han B., Fu XB., Han B., Lei YH., Chen W., Sun TZ., 2007, “*Chemotactic effects of burn rat serum on mesenchymal stem cells derived from different sources*”, *Zhonghua Shao Shang Za Zhi*; 23(1):25-8.
- Heo SC., Jeon ES., Lee IH., Kim HS., Kim MB., Kim JH., 2011, “*Tumor necrosis factor-alpha-activated human adipose tissue-derived mesenchymal stem cells accelerate cutaneous wound healing through paracrine mechanisms*”, *J. Invest. Dermatol.*, 1559–1567.
- Jiang B., Liao R., 2010, “*The paradoxical role of inflammation in cardiac repair and regeneration*”, *J. Cardiovasc. Transl. Res.*, 410–416.
- Kwon YW., Heo SC., Jeong GO., Yoon JW., Mo WM., Lee MJ., et al., 2013, “*Tumor necrosis factor- α -activated mesenchymal stem cells promote endothelial progenitor cell homing and angiogenesis*”, *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 2136–2144.
- Lee MJ., Kim J., Kim MY., Bae YS., Ryu SH., Lee TG., Kim JH., 2010, “*Proteomic analysis of tumor necrosis factor-alpha-induced secretome of human adipose tissue-derived mesenchymal stem cells*”, *J. Proteome Res.*, 1754–1762.
- Li L., Yang M., Wang C., Zhao Q., Liu J., Zhan C., Liu Z., Li X., Wang W., Yang X., 2012, “*Effects of cytokines and chemokines on migration of mesenchymal stem cells following spinal cord injury*”, *Neural Regen Res*; 7:1106-12.

- Locksley RM., Killeen N., Lenardo MJ., 2001, “*The TNF and TNF receptor superfamilies: integrating mammalian biology*”, *Cell*, 104; 487–501.
- Plebanski M, Proudfoot O, Pouniotis D. *et al.* 2002. *immunogenetics and the design of plasmodium falciparum vaccines for use in malaria-endemic populations.* *J.Clin. Invest.*295-301.
- Ponte AL., Marais E., Gallay N., Langonne A., Delorme B., Herault O., Charbord P., Domenech J., 2007, “*The in vitro migration capacity of human bone marrow mesenchymal stem cells: comparison of chemokine and growth factor chemotactic activities*”, *Stem Cells*; 25:1737.
- Rustad KC., Gurtner GC., 2012, “*Mesenchymal Stem Cells Home to Sites of Injury and Inflammation*”, *Adv Wound Care (New Rochelle)*; 1(4): 147–152.
- Rüster B., Göttig S., Ludwig RJ., et al., 2006, “*Mesenchymal stem cells display coordinated rolling and adhesion behavior on endothelial cells*”, *Blood*; 108(12):3938-3944.