

DAFTAR PUSTAKA

- Baraniak P.R., McDevitt T.C., 2010, Stem cell paracrine actions and tissue regeneration, *Regen Med.* 5(1): 121–143.
- Broughton I.G., Janis J.E., Attiger C.E., 2006, Wound healing: an overview, *Plast Reconstr Surg* 117 (suppl) : 1 eS-32eS
- Chen L., Tredget E.E., Wu P.Y., 2008, Paracrine factors of mesenchymal stem cells recruit macrophages and endothelial lineage cells and enhance wound healing, *PLoS One* 3:e1886.
- Crisostomo P.R., Wang Y., Markel T.A., Wang M., Lahm T., Meldrum D.R., 2008, Human mesenchymal stem cells stimulated by TNF-alpha, LPS, or hypoxia produce growth factors by an NF kappa B- but not JNK-dependent mechanism, *Am J Physiol Cell Physiol* 294(3):C675–C682.
- Dominici M., Le Blanc K., Mueller I., Slaper-Cortenbach I., Marini F., Krause D., Deans R., Keating A., Prockop Dj., Horwitz E., 2006, Minimal Criteria for Defining Multipotent Stromal Cells, *Cytotherapy* 8(4): 315 317
- Gnecchi M., Zhang Z., Ni A., 2008, Paracrine mechanisms in adult stem cell signaling and therapy, *Circ Res.* 103:1204–1219.
- Gurtner, G.C., 2007, Wound Healing : Normal and Abnormal in Plastic Surgery.6 th ed Philadelphia, Lippincott William dan Wilkin, .p. 15-22.
- Heo S.C., Jeon E.S., Lee I.H., Kim H.S., Kim M.B., 2011, Tumor necrosis factor alpha-activated human adipose tissue-derived mesenchymal stem cells accelerate cutaneous wound healing through paracrine mechanisms, *J Invest Dermatol* 131: 1559-1567.
- Kang S.K., Shin, I.S., Ko M.S., Jo J.Y., Ra J.C., 2012, Journey of Mesenchymal Stem Cells for Homing: Strategies to Enhance Efficacy and Safety of Stem Cell Therapy, *Stem Cells International* (2012).
- Kwon, Y.W., Heo S.C., Jeong G.O., Yoon J.W., Mo W.M., Lee M.J., Jang I.H., Kwon S.M., Lee J.S., Kim J.H., 2013. Tumor necrosis factor- α -activated mesenchymal stem cells promote endothelial progenitor cell homing and angiogenesis, *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 1832(12).
- Lee M.J., Kim J., Kim M.Y., Bae Y.S., Ryu S.H., 2010, Proteomic analysis of tumor necrosis factor-alpha-induced secretome of human adipose tissue-derived mesenchymal stem cells, *J Proteome Res* 9:1754-1762.

- Luttenberger, Thomas., Schmid-Kotsas A., Menke A., Siech M., Beger H., Adler G., Grünert A., Bachem M.G., 2000, Platelet-Derived Growth Factors Stimulate Proliferation and Extracellular Matrix Synthesis of Pancreatic Stellate Cells: Implications in Pathogenesis of Pancreas Fibrosis, *Lab Invest* 80(1):47-55.
- Madrigal M., Rao K.S., Riordan N.H., 2014, A review of therapeutic effects of mesenchymal stem cell secretions and induction of secretory modification by different culture methods. *Journal of Translational Medicine* 2014, 12:260.
- Nandy D., Mukhopadhyay D., 2011, Growth Factor Mediated Signaling in Pancreatic Pathogenesis, *Cancers* 3:841-871
- Wang M., Crisostomo P.R., Herring C., Meldrum K.K., Meldrum D.R., 2006, Human progenitor cells from bone marrow or adipose tissue produce PDGF, HGF, and IGF-I in response to TNF by a p38 MAPK-dependent mechanism, *Am J Physiol Regul Integr Comp Physiol* 291: R880-R884.
- Wang S., Mo M., Wang J., Sadia S., Shi B., Fu X., Yu L., Tredget E.E., Wu Y., 2017, Platelet-derived growth factor receptor beta identifies mesenchymal stem cells with enhanced engraftment to tissue injury and pro-angiogenic property, *Cell Mol Life Sci*.