

DAFTAR PUSTAKA

- Aini, N, et al. (2008), Karakteristik Biologis dan Diferensiasi Stem Cell: Fokus pada *Mesenchymal Stem Cell*, cdk,161.
- Alfonso Eirin and Lilach O Lerman, (2014). Mesenchymal stem cell treatment for chronic renal failure. *Stem Cell Research & Therapy*, <http://stemcellres.com/content/5/4/83>
- Bellomo, R., Kellum, J. A., & Ronco, C. (2012). Acute kidney injury. *The Lancet*, 380(9843), 756–766. [https://doi.org/10.1016/S0140-6736\(11\)61454-2](https://doi.org/10.1016/S0140-6736(11)61454-2)
- Bonventre, J., & Yang, L. (2011). Cellular pathophysiology of ischemic acute kidney injury. *J Clin Invest.*, 121(11), 4210–4221. <https://doi.org/10.1172/JCI45161.4210>
- Bieback K., Kern S., Klüter H., Eichler H. (2010). Critical parameters for the isolation of mesenchymal stem cells from umbilical cord blood. *Stem Cells*. 2010;22(4):625-34.
- DeVos J. et al.(2014). Intermediate-Term Graft Loss After Renal Transplantation is Associated With Both Donor-Specific Antibody and Acute Rejection. *Clinical And Translational Research*.97:5,
- Dominici, M., Le Blanc, K., Mueller, I., Slaper-Cortenbach, I., Marini, F., Krause, D., et al. (2006). Minimal Criteria for Defining Multipotent Mesenchymal Stromal Cell. The International Society for Cellular therapy Position Statement. *NCBI*.
- Ezquer, Fernando.,dkk, (2015). Proregenerative Microenvironment Triggered by Donor Mesenchymal Stem Cells Preserves Renal Function and Structure in Mice with Severe Diabetes Mellitus. *BioMed Research International*: Hindawi Publishing. Volume 2015
- Ferrara N.(2010). Pathways mediating VEGF-independent tumor angiogenesis. *Cytokine Growth Factor Rev* 2010; 21(1): 21-6.
- Halim, D., Murti, H., Sandra, F., Boediono, A., Djuwantono, T., & Setiawan, B. (2010). *Stem Cell: Dasar Teori dan Aplikasi Klinis*. Jakarta: Erlangga.
- Han YF, Tao R, Sun TJ, Chai JK, Xu G, Liu J. (2013): Optimization of human umbilical cord mesenchymal stem cell isolation and culture methods. *Cytotechnology*. 65(5):819–27. doi:10.1007/s10616-012-9528-0.

- Hou, J., Zhong, T., Guo, T., Miao, C., Zhou, C., Long, H., Wang, T. (2017). Apelin promotes mesenchymal stem cells survival and vascularization under hypoxic-ischemic condition in vitro involving the upregulation of vascular endothelial growth factor. *Experimental and Molecular Pathology*, 102(2), 203–209. <https://doi.org/10.1016/j.yexmp.2017.01.015>
- Hoeben A, Landy B, et al. (2008) Vascular endothelial growth factor and angiogenesis. *Pharmacol Rev* 2008; 56: 549-80.
- Johnson K., W. T. (2013). Vascular Endothelial Growth Factor and Angiogenesis in the Regulation of Cutanepous Wound Repair. *Wound Healing Society*. 647-661.
- Kang S. K., I. s. (2012). Journey of mesenchymal stem cells for homing:strategies to enhance efficancy and safety of stem cell therapy. *stem cells international* .
- Kementrian Kesehatan RI Pusat, (2017). Situasi Ginjal Kronis. *Data dan Informasi*
- Kwon Y. W. ,& S. C. (2013). Tumor necrosis factor- α -activated mesenchymal stem cells promote Tumor necrosis factor-. *Biochimica et Biophysica Acta* , 2136-2144.
- Nagaishi, Kanna,dkk,(2016). Mesenchymal stem cell therapy ameliorates diabetic nephropathy via the paracrine effect of renal trophic factors including exosomes. *Scientific Reports*:| 6:34842 | DOI: 10.1038/srep34842
- Peired AJ, Sisti A, Romagnani P. (2016): Mesenchymal Stem Cell-Based Therapy for Kidney Disease: A Review of Clinical Evidence. *Stem Cells Int.*: 4798639. doi:10.1155/2016/4798639.
- Prodjosudjadi, W., A. Suhardjono. (2009) End-Stage Renal Disease in Indonesia: *Treatment Development*. Ethn Dis.2009;19(suppl1):S1-33- S1-36
- Triyanti, K., Suhardjono, P. Soewondo, H. (2012). Shatri Renal Function Decrement in Type 2 Diabetes Mellitus Patient in Cipto Mangunkusumo Hospital. *Acta*
- Verma, S. K., & Molitoris, B. A. (2015). Renal Endothelial Injury and Microvascular Dysfunction in Acute Kidney Injury. *Seminars in Nephrology*, 35(1), 96–107. <https://doi.org/10.1016/j.semnephrol.2015.01.010>

Wang, M., et al. (2006). Human progenitor cells from bone marrow or adipose tissue produce VEGF, HGF, and IGF-I in response to TNF by a p38 MAPK-dependent mechanism. *Am J Physiol Regul Integr Comp Physiol* 291: R880–R884.

Wulandari,B & Ismail .A, cit Chin, (2008), Pengaruh Pemberian Seduhan Kelopak Bunga Rosella,(*hibiscus sabdarriffa*) dosis bertingkat selama 30 hari terhadap gambaran histologi Ginjal Tikus Wistar.

Yoshida S., et al. (2010). Involvement of interleukin-8, vascular endothelial growth factor, and basic fibroblast growth factorin tumor necrosis factor alpha-dependent angiogenesis. *Mol Cel Biol*; 17(7): 4015-23.