

DAFTAR PUSTAKA

- Ang, M, Chaurasia S., Angunawela R. (2018) ‘Femtosecond Lenticule Extraction (FLEEx): Clinical Results , Interface Evaluation , and Intraocular Pressure Variation AND’, 53, pp. 1414–1421. doi: 10.1167/iovs.11-8808.
- Bashir, Z. S., Ali M., Anwar. (2017) ‘Femto-LASIK: The recent innovation in laser assisted refractive surgery’, *Journal of the Pakistan Medical Association*, 67(4), pp. 609–615.
- Caesarya, S., Wangsaatmadja H., Rini M. (2015) ‘Clinical Outcomes of Laser In Situ Keratomileusis (LASIK) Using Microkeratome and Laser Femtosecond Flap in Myopic Patients’, *Ophthalmol Ina*, 41(2), pp. 165–170.
- Ekktet, C., Morakot T., Sukanda. (2015) ‘Efficacy, predictability, and safety of small incision lenticule extraction: 6-months prospective cohort study’, *BMC Ophthalmology*. Eye and Vision, 2. doi: 10.1186/s40662-015-0024-4.
- Ganesh, S. and Gupta, R. (2014) ‘Comparison of Visual and Refractive Outcomes Following Femtosecond Laser-Assisted LASIK With SMILE in Patients With Myopia or Myopic Astigmatism’, *Journal of Refractive Surgery*, 30(9), pp. 590–596. doi: 10.3928/1081597X-20140814-02.
- Gardiner, J. P., Moshirfar M., Schliesser J. (2010) ‘Laser in situ keratomileusis flap complications using mechanical microkeratome versus femtosecond laser : Retrospective comparison’, *Journal of Cataract & Refractive Surgery*. ASCRS and ESCRS, 36(11), pp. 1925–1933. doi: 10.1016/j.jcrs.2010.05.027.
- Holden, B. A., Fricke T., Wilson D. (2016) ‘Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050’, *Ophthalmology*. American Academy of Ophthalmology, 123(5), pp. 1036–1042. doi: 10.1016/j.ophtha.2016.01.006.
- Kaiser, P. K. (2009) ‘Prospective evaluation of visual acuity assessment: a comparison of snellen versus ETDRS charts in clinical practice (An AOS Thesis).’, *Transactions of the American Ophthalmological Society*, 107, pp. 311–24. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2814576/>&tool=pmcentrez&rendertype=abstract.
- Karim M., Riau A., Lwin N. *et al.* (2014) ‘Early Corneal Nerve Damage and Recovery Following Small Incision Lenticule Extraction (SMILE) and

Laser In'. doi: 10.1167/iovs.13-13324.

Lazaridis, A., Droutsas K., Sekundo W. (2017) 'Clinical Study Corneal Clarity and Visual Outcomes after Small-Incision Lenticule Extraction and Comparison to Femtosecond Laser-Assisted In Situ Keratomileusis', 2017.

Lim, C. H. L, Riau A., Lwin N. (2013) 'LASIK following small incision lenticule extraction (SMILE) lenticule re-implantation: A feasibility study of a novel method for treatment of presbyopia', *PLoS ONE*, 8(12), pp. 1–12. doi: 10.1371/journal.pone.0083046.

Lin, F., Xu, Y. and Yang, Y. (2014) 'Comparison of the Visual Results After Relex SMILE and Femtosecond LASIK'. doi: 10.3928/1081597X-20140320-03.

Morgan, I. G., Ohno-Matsui, K. and Saw, S. M. (2012) 'Myopia', *The Lancet*, 379(9827), pp. 1739–1748. doi: 10.1016/S0140-6736(12)60272-4.

Muñoz, G., Albarrán-diego, C. and Ferrer-blasco, T. (2010) 'Long-term comparison of corneal aberration changes after laser in situ keratomileusis : Mechanical microkeratome versus femtosecond laser flap creation', *Journal of Cataract & Refractive Surgery*. ASCRS and ESCRS, 36(11), pp. 1934–1944. doi: 10.1016/j.jcrs.2010.06.062.

Reilly, C. D., Panday V., Lazos V. (2010) 'PRK vs LASEK vs Epi-LASIK: a comparison of corneal haze, postoperative pain and visual recovery in moderate to high myopia.', *Nepalese journal of ophthalmology : a biannual peer-reviewed academic journal of the Nepal Ophthalmic Society : NEPJOPH*, 2(2), pp. 97–104. doi: 10.3126/Nepal Ophthalmic Society.v2i2.3715.

Riau, A. K., Angunewela R., Chaurasia S. (2011) 'Early Corneal Wound Healing and Inflammatory Responses after Refractive Lenticule Extraction (ReLEX)', 52(9). doi: 10.1167/iovs.11-7439.

Usman, S., Nukman, E. and Bebasari, E. (2014) 'Hubungan antara Faktor Keturunan, Aktivitas Melihat Dekat dan Sikap Pencegahan Mahasiswa Fakultas Kedokteran Universitas Riau terhadap Kejadian Miopia', *Jom Fk*, 1(2), pp. 1–13. Available at: <http://download.portalgaruda.org/article.php?article=186994&val=6449&title=HUBUNGAN+ANTARA+FAKTOR+KETURUNAN,+AKTIVITAS+MELIHAT+DEKAT+DAN+SIKAP+PENCEGAHAN+MAHASISWA+FAKULTAS+KEDOKTERAN+UNIVERSITAS+RIAU+TERHADAP+KEJADIAN+MIOPIA>.

- .Vestergaard, A., Ivarsen A., Sven A. (2013) ‘Femtosecond (FS) Laser Vision Correction Procedure for Moderate to High Myopia : a Prospective Study of ReLEx Ô flex and Comparison with a Retrospective Study of FS-Laser in Situ Keratomileusis’, *Achta Ophthalmology*, 91, pp. 355–362. doi: 10.1111/j.1755-3768.2012.02406.x.
- Widodo, A. and Prillia, T. (2007) ‘Miopia Patologi’, *Jurnal Oftalmologi Indonesia*, 5(1), pp. 19–26. Available at: <http://repository.usu.ac.id/bitstream/123456789/44769/4/Chapter II.pdf>.
- Yanoff, M. and Duker, jay s. (2014) *Ophthalmology*. fourth edi. ELSEVIER SAUNDERS.
- Yu, L. Li Z., Gao J. (2011) ‘Epidemiology, genetics and treatments for myopia.’, *International journal of ophthalmology*, 4(6), pp. 658–69. doi: 10.3980/j.issn.2222-3959.2011.06.17.
- Zhang, J., Zhou Y., Zheng Y. (2017) ‘Comparison of visual performance recovery after thin-flap LASIK with 4 femtosecond lasers’, *International journal of ophthalmology*, 10(10), p. 1566. doi: 10.18240/ijo.2017.10.14.