

DAFTAR PUSTAKA

- Acarturk, E., Cayli, M., Akpınar, O., Attila, G., Demir, M., 2004, Relation between age and gender differences in plasma triglyceride concentrations and coronary artery disease in Southern Turkey, *International Journal of Clinical Chemistry*, p:123.
- Adam, J.M.F, 2009, Dislipidemia dalam *Buku Ajar Ilmu Penyakit Dalam*, Edisi 5, cetakan I, editor edisi bahasa Indonesia: Aru W Sudono dkk, FKUI, Jakarta, p:1984-1986.
- Adebayo, J.O., Akinyinka, A.O., Odewole, G.A., Okwusidi, J.I., 2007, Effect of Caffeine on the Risk of Coronary Heart Disease a Re-Evaluation, *Indian Journal of Clinical Biochemistry*, 22(1),p:29,30.
- ALboqai, O.K., Saraireh, R.M., Saif, M.A., Diab, M.M., 2006, Generalized Obesity, Serum Lipid and Lipoproteins in Adult Males in a Semi Urban Community in The North of the Jordan, *JRMS*, 13(1), p:39-45.
- Angulo, P., Lindor, K, D., 2002, Non-alcoholic fatty liver disease, *Journal of Gastroenterology and Hepatology*, p: 186-190.
- Apovian, C.M., 2004, Sugar-Sweetened Soft Drinks, Obesity, and Type 2 Diabetes, *JAMA*, p: 978.
- Assy, Nimer.Nasser., Gattas, Kamayse., Iad, Nseir., William, Beniashvili., Zaza, Djibre., Agness, Grosovski, Maria., 2008, Soft Drink Consumption Linked With Fatty Liver In The Absence Of Traditional Risk Factors, *Can J Gastroenterol*, 22, p: 811.
- Australian Beverage Council, 2013,
www.australianbeverages.org/products/products-carbonated-regular-and-diet-soft-drinks, *Carbonated soft drinks* , Di kutip tanggal 22 Mei 2013.
- Beigh,S,H., Jain,S., 2012, Prevalence of metabolic syndrome and gender differences, *Biomedical Informatics*, p:613.
- Botezelli, J.D, Dalia, R.A., Reis, I.M., Barbieri, R.A., Rezende, T.M., Pelarigo, J.G., Codogno, J., Goncalves, R., Mello, M.A., 2010, Chronic Consumption of Fructose Rich Soft Drinks Alters Tissue Lipids of Rats, *dmsjournal*, 2,p:1-8.
- Botham, K.M., Mayes, P.A., 2009, Pengangkutan dan Penyimpanan Lipid dalam *Biokimia Harper*, Edisi 27, cetakan I, editor edisi bahasa Indonesia: Nanda Wulandari dkk, EGC, Jakarta, p: 234.

- Bray, G.A., Nielsen, S.J., Popkin, B.M., 2004, Consumption of high-fructose corn syrup in beverages may play a role in the epidemic of obesity, *The American Journal of Clinical Nutrition*, 79, p:537.
- British Soft Drinks Association, 2013, www.britishsoftdrinks.com, *Ingredients of Soft Drinks*, Di kutip tanggal 22 Mei 2013.
- Caceaune, E., 2012, The Relationship Between Nonalcoholic Fatty Liver Disease and The Metabolic Syndrome, *the Romanian Academy*, p : 143-150.
- Champe, P.C., Harvey, R.A., Ferrier, D.R., 2004, Biokimia Ulasan Bergambar, *Metabolisme Lipid*, Edisi 3, Alih bahasa: Novrianti A dkk, EGC, Jakarta.
- Choi, J.W., Pai, S.H., Kim, S.K., 2002, Associations Between Total Body Fat and Serum Lipid Concentrations in Obese Human Adolescents, *Annals of Clinical & Laboratory Science*, p: 271-277.
- Dahlan, M. S., 2008. Statistik Untuk Kedokteran dan Kesehatan, edisi 3, Salemba Medika: Jakarta. p: 55, 63, 66, 89.
- Dhingra, Ravi., Sullivan, Lisa., Jacques, P.F., Wang, T.J., Fox, C.S., Meigs, J.B., D'Agostino, R.B., Gaziano, J.M., Vasan, R.S., 2007, Soft Drink Consumption and Risk of Developing Cardiometabolic Risk Factors and the Metabolic Syndrome in Middle-Aged Adults in the Community, *Journal of the American Heart Association*, 116, p: 483.
- Dilapanga, Alfira, 2008, Faktor-Faktor Yang Berhubungan Dengan Perilaku Konsumsi Soft Drinks Pada Siswa Smp Negeri 1 Ciputat Tahun 2008, Program Studi Kesehatan Masyarakat Fakultas Kedokteran dan Ilmu Kesehatan Universitas Islam Negeri Syarif Hidayatullah, Jakarta.
- Ducommun, D., 2007, <http://www.rattycorner.com/odds/age.shtml>, *Ratty Corner and the Brandywine Rats*, Dikutip tanggal 23 Agustus 2013.
- El-Ghany, A., Rasha, N.M., Nagar, El-Saiyed, M., 2012, Hypolipidemic Effect of Caffeine Beverages in Fatty Liver Injured Rats, *Journal of Applied Sciences Research*, 8(3), p: 1502.
- Emanuele, N.V., Swade, T.F, Emanuele, N.A., 1998, Consequences of Alcohol Use in Diabetics, *Alcohol Health & Research World*, p: 216.
- Fredhlo, B.B., Battig, K., Holmet, J., Nehlig, A., Zvartau, E.E., 1999, Actions of Caffeine in the Brain with Special Reference to Factors That Contribute to Its Widespread Use, *The American Society for Pharmacology and Experimental Therapeutics*, 51, p:86.
- Gaby, A.R., 2005, Adverse Effects of Dietary Fructose, *Alternative Medicine Review*, 10, p: 294.

- Ghezzi, A.C., Cambri L.T., Ribeiro, C., Botezelli, J.D., Mello, M.A.R., 2011, Impact of Early Fructose Intake on Metabolic Profile And Aerobic Capacity of Rats, *Lipids in Health and Disease*, 10,p:1.
- Guyton, A.C., Hall, J.E., 2007, Metabolisme Lipid dalam *Buku Ajar Fisiologi Kedokteran*, edisi 11, editor bahasa Indonesia: Luqman Yanuar Rachman dkk, Alih Bahasa: Irawati dkk, EGC, Jakarta, p: 882-889.
- Hostmark, T.A., Lunde, H.S.M., Eilersten, E., 2010, Does an acid load promote liver desaturases and increase serum lipids?, *Academic Journals*, p:008-016.
- Hu, F.B., 2010, Sugar Sweetened Beverages as a Major Modifiable Risk Factor for type 2 Diabetes, *CMR journal*, p: 9.
- Khalil, S.M.R., 1999, Some Biochemical Studies On Serum Lipids Content In Rats Before And After Caffeine Administration, Zagazig University, p:28.
- Kobayashi-Hattori, K., Mogi, A, Matsumoto, Y., Takita, T., 2005, Effect Caffeine on the Body Fat and Lipid Metabolism of Rats Fed on a High-Fat Diet, *JSBA*, 69(11),p: 2219.
- Kotronen, A., Yki-Jarvinen, H., 2008, Fatty Liver: A Novel Component of the Metabolic Syndrome, *Journal of the American Heart Association*, p : 28-30.
- Kuk, J.L., Ardern, C.I., 2010, Age and Sex Differences in the Clustering of Metabolic Syndrome Factors, *diabetesjournals*, p:2437-2460.
- Kusumawati, D., 2004, Bersahabat Dengan Hewan Coba, Gajah Mada University Press, Yogyakarta, p: 8-9.
- Lane, J.D., Pieper, C.F., Phillips-Bute, B.G., Bryant, J.E., Kuhn, C.M., 2002, Caffeine Affects Cardiovascular and Neuroendocrine Activation at Work and Home, *The American Psychosomatic Medicine*, 64, p:595.
- Malik, V.S., Schulze, M.B., Hu, F.B., 2006, intake of sugar-sweetened beverages and weight gain: a systematic review, *American Journal of Clinical Nutrition*, 84, p:275.
- Parks, E.J., Krauss, R.M., Christiansen, M.P., Neese, R.A., Hellerstein, M.K., 1999, Effects of a Low-Fat, High-Carbohydrate Diet on VLDL-Triglyceride Assembly, Production, And Clearance, *The Journal of Clinical Investigation*, 104, p:1087.
- Prahastuti, S., 2011, Konsumsi Fruktosa Berlebihan dapat Berdampak Buruk bagi Kesehatan Manusia, Fakultas Kedokteran, Universitas Kristen Maranatha Bandung 10, p: 173-189.

- Ree, M., Erik, S., 2011, Simultaneous determination of aspartame, benzoic acid, caffeine, and saccharin in sugar-free beverages using HPLC, *Concordia College Journal of Analytical Chemistry I*, p : 73-77.
- Ryu, MH., Cha,YS., 2003, The Effects of a High-fat or High-sucrose Diet on Serum Lipid Profiles, Hepatic Acyl-CoA Synthetase, Carnitine Palmitoyltransferase-I, and the Acetyl-CoA Carboxylase mRNA Levels in Rats, *Journal of Biochemistry and Molecular Biology*, 36, p:312-314.
- Schoppen,S., Perez-Granadoz,A,M., Carbajal,A., Sarria,B., Sanchez-Muniz,F,J., 2005, Sodium Bicarbonated Mineral Water Decreases Postprandial Lipaemia in Postmenopausal Women Compared to a Low Mineral Water, *British Journal of Nutrition*, p:582-587.
- Sherwood, L., 2001, Atherosclerosis: Cholesterol and Beyond dalam Human Physiology: From Cells to Systems, *Nelson Education, Ltd.*, Canada.
- Stanhope, K.L., Schwarz, J.M., Keim, N.L., Griffen, S.C., Bremer, A.A., J.L., Graham, B., Hatcher, C.L., Cox, A., Dyachenko, W., Zhang, J.P., McGahan, A., Seibert, R.M., Krauss, S., Chiu, E.J., Schaefer, M., Ai, S., Otokozawa, K., Nakajima, T., Nakano, C., Beysen, M.K., Hellerstein, L., Berglund, P.J., Havel, 2009, Consuming Fructose-Sweetened, Not Glucose Sweetened, Beverages Increases Visceral Adiposity and Lipids and Decreases Insulin Sensitivity In Overweight/Obese Humans, *The Journal of Clinical Investigation*, 119, p: 1323.
- Stanhope, K.L., Havel, P.J., 2008, Fructose consumption: potential mechanisms for its effects to increase visceral adiposity and induce dyslipidemia and insulin resistance, p: 20.
- Supriyono, M., 2008, Faktor-Faktor Risiko Yang Berpengaruh Terhadap Kejadian Penyakit Jantung Koroner Pada Kelompok Usia < 45 Tahun Studi Kasus Di RSUP dr. Kariadi dan RS Telogorejo UNDIP, Semarang, p:1-94.
- Tsalissavrina, Iva., Wahono Djoko, Handayani Dian, 2006, Pengaruh Pemberian Diet Tinggi Karbohidrat Dibandingkan Diet Tinggi Lemak Terhadap Kadar Trigliserida dan HDL Darah Pada Rattus Novergicus Galur Wistar, *Jurnal Kedokteran Brawijaya*, 22, p: 87.
- WHO, World Health Organization, 1993, Research Guidelines for the Safety Effecacy of Herbal Medicine, Regional Office for The Watern Pacific, Manila.
- Yule, A., 2002, Increased Soft Drink Consumption is contributing to an Increased Incidence of Obesity, *nutrition byte*, University of California Los Angeles, p: 2.
- Zair, Y., Kasbi-Chadli, F., Housez, B., Pichelin, M., Cazaubiel, M., Raoux, F., 2013, Effect Of a High Bicarbonate Mineral Water on Fasting and Postprandial Lipemia in Moderately Hypercholesterolemic Subjects: a Pilot Study, *Biomed Central*, p: 2-7.