

DAFTAR PUSTAKA

- Abdo, R. A. *et al.* (2019) ‘Prevalence and contributing factors of birth asphyxia among the neonates delivered at Nigist Eleni Mohammed memorial teaching hospital, Southern Ethiopia: A cross-sectional study’, *BMC Pregnancy and Childbirth*, 19(1), pp. 1–7. doi: 10.1186/s12884-019-2696-6.
- ACOG (2020) ‘ACOG Practice Bulletin Clinical Management Guidelines for Obstetrician Gynecologists. Gestational Hypertension and Preeclampsia’, *Obstetrics & Gynecology*, 135(6), pp. e237–e260.
- Aslam, H. M. uhamma. *et al.* (2014) ““Risk factors of birth asphyxia””, *Italian journal of pediatrics*, 40, p. 94. doi: 10.1186/s13052-014-0094-2.
- B, A. N. *et al.* (2018) ‘A Retrospective Study on the Risk of Respiratory Distress Syndrome in Singleton Pregnancies with Preterm Premature Rupture of Membranes between 24 + 0 and 36 + 6 Weeks , Using Regression Analysis for Various Factors’, 2018, pp. 1–7.
- Bedie, N. A., Wodajo, L. T. and Mengesha, S. T. (2019) ‘Magnitude and Determinants of Birth Asphyxia : Unmatched Case Control Study Assela Referral Teaching Hospital , Arsi Zone , Ethiopia’, 7(1). doi: 10.19080/GJORM.2019.07.555705.
- BKKBN, BPS and Kemenkes RI (2018) ‘Survei Demografi Kesehatan Indonesia’, *Usaid*, pp. 1–606.
- Cunningham, F. G. *et al.* (2018) *Williams Obstetric*. 25 th. United States: Mc Graw Hills.
- damayanti, mutia (2017) ‘Hubungan Preeklamsia dengan Kejadian Asfiksia Neonatorum pada bayi baru lahir’.
- Gupta, Shruti, Malik, S. and Gupta, Shailesh (2020) ‘Neonatal complications in women with premature rupture of membranes (PROM) at term and near term and its correlation with time lapsed since PROM to delivery’, *Tropical Doctor*, 50(1), pp. 8–11. doi: 10.1177/0049475519886447.
- Harrison, S. (no date) *No Title*.
- Health, P. (2020) ‘Risk Factors of Perinatal Asphyxia Among Newborns Delivered at Public Hospitals in Addis Ababa , Ethiopia : Case – Control Study’, pp. 297–306.
- Ibrahim, N. A., Muhye, A. and Abdulie, S. (2017) ‘Prevalence of Birth Asphyxia and Associated Factors among Neonates Delivered in Dilchora Referral Hospital, in Dire Dawa, Eastern Ethiopia’, *Clinics in Mother and Child Health*, 14(4). doi: 10.4172/2090-7214.1000279.

- Id, A. W., Muluneh, A. and Denis, C. S. (2019) ‘Birth asphyxia and its associated factors among newborns in public hospital , northeast’, pp. 1–13.
- Kalteren, W. S. *et al.* (2018) ‘Perinatal Anemia is Associated with Neonatal and Neurodevelopmental Outcomes in Infants with Moderate to Severe Perinatal Asphyxia’, *Neonatology*, 114(4), pp. 315–322. doi: 10.1159/000490369.
- Kardana, I. M. (2016) ‘Risk Factors of Perinatal Asphyxia in The Term Newborn at Sanglah General Hospital, Bali-Indonesia’, *Bali Medical Journal*, 5(1), p. 196. doi: 10.15562/bmj.v5i1.312.
- Kemenkes (2020) ‘Pokok-Pokok Renstra Kemenkes 2020-2024’, *Pokja Renstra Kemenkes 2020-2024*, pp. 1–40.
- Kesehatan, D. and Jawa, P. (2019) ‘DINAS KESEHATAN PROVINSI JAWA TENGAH DINAS KESEHATAN PROVINSI JAWA TENGAH’.
- Klinger, G. *et al.* (2020) ‘Antepartum Hemorrhage and Outcome of Very Low Birth Weight, Very Preterm Infants: A Population-Based Study’, *American Journal of Perinatology*, 1(212). doi: 10.1055/s-0040-1710353.
- Kurniati, D. P. Y. (2019) ‘Determinant Factors of Asphyxia Neonatorum Risk in Neonates in Regional General Hospitals Biak Numfor District’, *Galore International Jurnal of Applied Sciences and Humanities*, 3(1), pp. 59–69.
- Kusumaningrum, R. Y., Murti, B. and Prasetya, H. (2019) ‘Low Birth, Prematurity, and Pre-Eclampsia as Risk Factors of Neonatal Asphyxia’, *Journal of Maternal and Child Health*, 04(01), pp. 49–54. doi: 10.26911/thejmch.2019.04.01.07.
- Maulida, H. (2019) ‘Asfiksia Neonatorum di RS PKU Muhammadiyah Bantul’.
- Moshiro, R., Mdoe, P. and Perlman, J. M. (2019) ‘A Global View of Neonatal Asphyxia and Resuscitation’, *Frontiers in Pediatrics*, 7(November), pp. 1–6. doi: 10.3389/fped.2019.00489.
- Mulia, S. and Conference, I. (2017) ‘Advances in Health Science Research, volume 6 2nd Sari Mulia International Conference on Health and Sciences (SMICHS 2017)’, 6(Smichs), pp. 12–20.
- Mushtaq, R., Abbas, A. and Ahmed, W. (2019) ‘Perinatal outcomes in pregnant patients presenting with antepartum hemorrhage: Our experiences at a teaching hospital in Central Punjab.’, *The Professional Medical Journal*, 26(10), pp. 1645–1650. doi: 10.29309/tpmj/2019.26.10.3152.
- Opitasari, C. and Andayasaki, L. (2016) ‘Maternal education, prematurity and the risk of birth asphyxia in selected hospitals in Jakarta’, *Health Science Journal of Indonesia*, 6(2), pp. 111–115. doi: 10.22435/hsji.v6i2.4769.111-115.
- Polglase, G. R. *et al.* (2017) ‘birth asphyxia’, 43(3), pp. 469–483. doi:

10.1016/j.clp.2016.04.006.Cardiovascular.

Rainaldi, M. A. (2016) ‘Pathophysiology of Birth Asphyxia’, *Clinics in Perinatology*, 43(3), pp. 409–422. doi: 10.1016/j.clp.2016.04.002.

Rana, S. et al. (2019) ‘Compendium on the Pathophysiology and Treatment of Hypertension’, pp. 1094–1112. doi: 10.1161/CIRCRESAHA.118.313276.

Restu, S. et al. (2017) ‘Relationship of Chronic Energy Deficiency in Pregnant Women with Low Birth Weight Newborn in Central Sulawesi Province’, *International Journal of Sciences: Basic and Applied Research*, 36(2), pp. 252–259.

Statement, P. (2020) ‘The Apgar Score’. doi: 10.1542/peds.2006-0325.

Tasew, H. et al. (2018) ‘Risk factors of birth asphyxia among newborns in public hospitals of Central Zone, Tigray, Ethiopia 2018’, *BMC Research Notes*, 11(1), pp. 1–7. doi: 10.1186/s13104-018-3611-3.

UNICEF Indonesia (2015) ‘Country Profile of Indonesia’, (000), pp. 142–143.

Vc, P. et al. (2020) ‘Yilgwan CS CC – BY Comparing neonatal outcomes in women with preeclampsia and those with normal pregnancy’, 47(3), pp. 258–263.

WHO (2020) ‘Newborns: Improving survival and well-being’.

Wosenu, L. et al. (2018) ‘Determinants of birth asphyxia among live birth newborns in University of Gondar referral hospital, northwest Ethiopia: A case-control study’, *PLoS ONE*, 13(9), pp. 1–12. doi: 10.1371/journal.pone.0203763.

Wu, C., Chen, C. and Chou, H. (2017) ‘ScienceDirect Pulmonary Hypoplasia Induced by Oligohydramnios : Findings from Animal Models and a Population-Based Study’, *Pediatrics and Neonatology*, 58(1), pp. 3–7. doi: 10.1016/j.pedneo.2016.04.001.

Yadav, N. and Damke, S. (2017) ‘Study of risk factors in children with birth asphyxia’, *International Journal of Contemporary Pediatrics*, 4(2), p. 518. doi: 10.18203/2349-3291.ijcp20170701.