

## DAFTAR PUSTAKA

- Amano, O. *et al.* (2012) 'Anatomy and histology of rodent and human major salivary glands: Overview of the Japan salivary gland society-sponsored workshop', *Acta Histochemica et Cytochemica*, 45(5), pp. 241–250. doi: 10.1267/ahc.12013.
- Anindita, Y. P. C., Raymondalexas, M. C. and Suryo, P. Y. (2010) 'Hubungan antara Pemberian Radioterapi dengan Terjadinya Distress , Anxiety dan Depresi pada Penderita Kanker Payudara', *Berita Kedokteran Masyarakat*, 26(1), pp. 1–6. doi: 10.1007/978-3-319-12817-7.
- Arpa, S. and Jubhari, E. H. (2017) 'Sifat saliva dan hubungannya dengan pemakaian gigi tiruan lepasan', *Makassar Dental Journal*, 6(2), pp. 78–82. Available at: <http://jurnal.pdgimakassar.org/index.php/MDJ/article/view/26/25>.
- Arrifin, A. *et al.* (2018) 'The Effect of Radiotherapy for Treatment of Head and Neck Cancer on Oral Flora and Saliva.', *Oral health & preventive dentistry*, 16(5), pp. 425–429. doi: 10.3290/j.ohpd.a41364.
- Chitapanarux, I. and Iamaroon, A. (2020) 'Salivary glands and dental complications after radiotherapy for nasopharyngeal carcinoma', *Annals of Nasopharynx Cancer*, 4, pp. 7–7. doi: 10.21037/anpc-20-17.
- Devi, B. C., Yueniwati, Y. and Dw, A. (2016) 'Comparison of Caspase-3 Responses of Tracheal Cells to Gamma Rays Radiation in Single Dose and Fractination Dose', 3(3), pp. 121–127.
- Dwikuntari, L., Setijadi, A. R. and Hendrik (2017) 'External beam radiation therapy pada kanker paru', *Berkala Ilmiah Kedokteran Duta Wacana*, 02(02), pp. 375–392. doi: 10.1007/978-1-4939-1145-5\_26.
- F., P. and Waschke, J. (2013) *Sobotta Atlas Anatomi Manusia :Jilid 3, Indo American Journal of Pharmaceutical Sciences*. Jakarta: EGC. doi: 10.5281/zenodo.1477753.
- Fang, F. M. *et al.* (2010) 'Multivariate analysis of quality of life outcome for nasopharyngeal carcinoma patients after treatment', *Radiotherapy and Oncology*, 97(2), pp. 263–269. doi: 10.1016/j.radonc.2010.05.022.
- Fithrony, M. T. (2012) *PENGARUH RADIOTERAPI AREA KEPALA DAN LEHER TERHADAP CURAH SALIVA UNIVERSITAS DIPONEGORO TAHUN 2012 Lembar Pengesahan Laporan Akhir Hasil Penelitian*.
- Fitriatuzzakiyyah, N., Sinuraya, R. K. and Puspitasari, I. M. (2017) 'Cancer Therapy with Radiation: The Basic Concept of Radiotherapy and Its Development in Indonesia', *Indonesian Journal of Clinical Pharmacy*, 6(4), pp. 311–320. doi: 10.15416/ijcp.2017.6.4.311.
- Helmerhorst, E. J. (2012) 'Anatomy and Physiology of Salivary Glands', *Saliva*

and oral health, pp. 115–134. doi: 10.1038/sj.bdj.2013.421.

Humphrey, S. P. and Williamson, R. T. (2001) 'A review of saliva: Normal composition, flow, and function', 85(2).

Indriana, T. (2010) 'The relationship between salivary flow rate and calcium ion secretion in saliva.', *Stomatognathic (J.K.G Unej)*, 7(2), pp. 29–31.

Kasuma, N. (2015) *Fisiologi dan Patologi Saliva. I, Andalas University Press. I. Padang: Andalas University Press. Available at: <http://repo.unand.ac.id/3650/1/01.Buku-Fisiologi-dan-Patologi-Saliva.pdf>.*

Komite Penanggulangan Kanker Nasional Kemenkes RI (2015) 'Panduan Penatalaksanaan Kanker Nasofaring', *Kementerian Kesehatan Republik Indonesia. Komite Penanggulangan Kanker Nasional.*, pp. 1–56.

Lal, P. *et al.* (2010) 'Changes in salivary flow rates in head and neck cancer after chemoradiotherapy', *Journal of Cancer Research and Therapeutics*, 6(4), pp. 458–462. doi: 10.4103/0973-1482.77105.

Larasati, A. (2016) 'PERBEDAAN DERAJAT KEASAMAN ( pH ) SALIVA OLEH':

Lee, N. *et al.* (2000) 'Nasopharyngeal Carcinoma', *Clinical Radiation Oncology. 3rd ed.*, pp. 618 – 38.

Liang, X. *et al.* (2016) 'Radiation caries in nasopharyngeal carcinoma patients after intensity-modulated radiation therapy: A cross-sectional study', *Journal of Dental Sciences*, 11(1), pp. 1–7. doi: 10.1016/j.jds.2015.09.003.

Lin, C. Y. *et al.* (2015) 'Effects of radiotherapy on salivary gland function in patients with head and neck cancers', *Journal of Dental Sciences*, 10(3), pp. 253–262. doi: 10.1016/j.jds.2015.01.004.

Marucci, L. *et al.* (2010) 'INFLUENCE OF INTENSITY-MODULATED RADIATION THERAPY TECHNIQUE ON XEROSTOMIA AND RELATED QUALITY OF LIFE IN PATIENTS TREATED WITH INTENSITY-MODULATED RADIATION THERAPY FOR NASOPHARYNGEAL CANCER', *Head and Neck*, 36(10), p. 1391. doi: 10.1002/HED.

Meka Anggidian Primadina1, M. I. (2017) 'Tumor Nasofaring dengan Diploopia Pada Pasien Usia 44 Tahun', *Jurnal Medula*, 7(4), pp. 181–186.

Pow, E. H. N. *et al.* (2006) 'Xerostomia and quality of life after intensity-modulated radiotherapy vs. conventional radiotherapy for early-stage nasopharyngeal carcinoma: Initial report on a randomized controlled clinical trial', *International Journal of Radiation Oncology Biology Physics*, 66(4), pp. 981–991. doi: 10.1016/j.ijrobp.2006.06.013.

Razek, A., Khalek, A. A. and King, A. (2012) 'MRI and CT of nasopharyngeal carcinoma', *American Journal of Roentgenology*, 198(1), pp. 11–18. doi: 10.2214/AJR.11.6954.

- Setyawan, A. and Djakaria, H. M. (2014) 'EFEK DASAR RADIASI PADA JARINGAN', *Journal of the Indonesian Radiation Oncology Society*, 5(1), pp. 1–41.
- Sim, C. P. C. *et al.* (2018) 'Xerostomia, salivary characteristics and gland volumes following intensity-modulated radiotherapy for nasopharyngeal carcinoma: a two-year follow up', *Australian Dental Journal*, 63(2), pp. 217–223. doi: 10.1111/adj.12608.
- Su, S. F. *et al.* (2012) 'Long-term outcomes of early-stage nasopharyngeal carcinoma patients treated with intensity-modulated radiotherapy alone', *International Journal of Radiation Oncology Biology Physics*, 82(1), pp. 327–333. doi: 10.1016/j.ijrobp.2010.09.011.
- Surjadi, N. and Amtha, R. (2013) 'Radiotherapy Reduced Salivary Flow Rate and Might Induced *C. albicans* Infection', *Journal of Dentistry Indonesia*, 19(1), pp. 1–6. doi: 10.14693/jdi.v19i1.124.
- Tamin, S. and Yassi, D. (2011) 'Penyakit kelenjar saliva dan peran sialoendoskopi untuk diagnostik dan terapi', 41(2), pp. 95–104.
- Vining, K. H. (2017) 'Anatomy, biogenesis, and regeneration of salivary glands', *NIH Public Access*, 24(May 2014). doi: 10.1159/000358776.
- Wu, V. W. C. *et al.* (2020) 'A longitudinal study on parotid and submandibular gland changes assessed by magnetic resonance imaging and ultrasonography in post-radiotherapy nasopharyngeal cancer patients', *BJR/Open*, 2(1), p. 20200003. doi: 10.1259/bjro.20200003.
- Wu, V. W. C., Ying, M. T. C. and Kwong, D. L. W. (2011) 'Evaluation of radiation-induced changes to parotid glands following conventional radiotherapy in patients with nasopharyngeal carcinoma', *British Journal of Radiology*, 84(1005), pp. 843–849. doi: 10.1259/bjr/55873561.
- Xu, J. *et al.* (2014) 'Effectiveness of oxygen nebulization at preventing radiotherapy-induced mucositis in patients with nasopharyngeal cancer', *International Journal of Nursing Sciences*, 1(2), pp. 176–179. doi: 10.1016/j.ijnss.2014.05.017.