

15. SITI THOMAS_Tender Coconut Water can Inhibits Inflamation due to Cigarette Smoke

by Siti Thomas

Submission date: 08-Jan-2022 04:14PM (UTC+0800)

Submission ID: 1738795374

File name: oconut_Water_can_Inhibits_Inflamation_due_to_Cigarette_Smoke.doc (407K)

Word count: 4532

Character count: 24806

Tender Coconut Water can Inhibits Inflammation due to Cigarette Smoke

Siti Thomas Zulaikhah^{1*}, Sampurna², Joko Wahyu Wibowo¹, Helmia Fitri Nurul Aini³, Arrizki Azka Pratama⁴

¹Department of Public Health, Faculty of Medicine, Universitas Islam Sultan Agung, Semarang, Indonesia

²Department of Clinical Pathology, Faculty of Medicine, Universitas Islam Sultan Agung, Semarang, Indonesia

³ Internship Doctor, Semarang, Indonesia

⁴Program Master of Biomedical Science Faculty of Medicine, Universitas Islam Sultan Agung, Semarang, Indonesia

Abstract: Cigarette smoke contains numerous free radicals that can trigger inflammatory response. The response stimulates the release of pro-inflammatory cytokines such as C-reactive protein (CRP), tumor necrosis factor-alpha (TNF- α), interleukin (IL)-6 and IL-8. Tender coconut water containing L-arginine has been shown to reduce the levels of inflammatory cytokines including TNF- α and IL-6. This study aims to determine the potential of tender coconut water in inhibiting inflammation in mice exposed to cigarette smoke. In this posttest only control group study, twenty-four male Wistar rats were randomly divided into 4 groups: K1 (standard *et libitum*+ aqua diet); K2 (standard diet *et libitum* + aqua + exposure to cigarette smoke 3 sticks/day); K3 (standard diet *et libitum* + aqua+ exposure to cigarette smoke 3 sticks/day + vitamin E at a dose of 1.8 mg/200 grBW/day); K4 (standard diet *et libitum* + aqua + exposure to cigarette smoke 3 sticks/day + tender coconut water at a dose of 8mL/200 grBW/day). The treatment was given for 14 days. On day 15, the blood samples were collected for to the evaluation of CRP, IL-6, and TNF- α using ELISA (enzyme-linked immunosorbent assay). The data were analyzed using one way ANOVA test. The results showed that the level of CRP, IL-6 and TNF α in the group administrated with tender coconut water was significantly lower compared to that of exposed to cigarette smoke alone ($p<0.05$). This study demonstrated that tender coconut water can inhibits inflammation in mice due to cigarette smoke.

Keywords: Tender coconut water, cigarette smoke, CRP, IL-6, TNF- α .

Introduction

Smoking is a risk factor for a wide range of diseases, disability mortality [1]. Half a million Americans died from smoking or exposure to secondhand smoke at early age, while another 16 million suffer from serious illnesses each year [2]. The latest statistical data according to the World Health Organization (WHO) shows the percentage of smokers in Indonesia accounts for 33.8%, with 21.37% deaths from smoking [3]. Cigarette smoke can cause chronic obstructive pulmonary disease (COPD), cardiovascular diseases and other diseases. Smoking have been shown to increase the risks of chronic systemic diseases through inflammatory mechanisms [4].

Cigarette smoke is a source of exogenous free radicals with more than 7000 chemicals. One cigarette contains 107 oxidant molecules [5]. There are two main phases in cigarette smoke that play a role in the pathogenesis namely the tar and gas phase. Both phases are rich in free radicals and non-free radicals. One puff

of cigarette smoke contains 1017 free radicals in the tar phase and 1015 in the gas phase. Nicotine found in cigarettes can increase ROS reactive oxygen species (ROS), decrease in the antioxidant capacity of enzymes and induce the formation of active compounds that interfere with blood flow by inducing inflammation in blood vessels [1].

Cigarette smoke exposure can cause airway inflammation characterized by an increase in macrophages, neutrophils and T lymphocytes. This triggers the release of pro-inflammatory cytokines such as tumor necrosis factor-alpha (TNF- α), interleukin (IL)-6 and IL-8 [6]. Several studies have shown increased levels of C-reactive protein (CRP), TNF-, IL-6 and IL-8 in COPD patients with inflammation due to exposure to cigarette smoke. Levels of C-reactive protein (CRP), TNF-, IL-6 and IL-8 in smoking COPD patients have been shown to be higher than that of non-smoking, exposure to cigarette smoke increases levels of pro-inflammatory

Received (date):

About the author: Siti Thomas Zulaikhah, Joko Wahyu Wibowo, Department of Public Health, Faculty of Medicine, Universitas Islam Sultan Agung, Semarang, Indonesia; Sampurna, Department of Pathological Clinic, Faculty of Medicine, Universitas Islam Sultan Agung, Semarang, Indonesia; Helmia Fitri Nurul Aini, Internship Doctor, Semarang, Indonesia; Arrizki Azka Pratama, Program Master of Biomedical Science, Faculty of Medicine, Universitas Islam Sultan Agung, Semarang, Indonesia

*Corresponding Author Siti Thomas Zulaikhah : sitithomas@unissula.ac.id

cytokines [5].

Tender coconut water contains L-arginine, polyphenols, vitamin C, selenium and minerals (Cu, Mg, Mn, K, Na, Zn). L-arginine in tender coconut water can significantly reduce free radicals. Oxidative stress decreases along with the decrease in free radicals in the body. L-arginine can inhibit inflammation indicated by the decrease in levels of pro-inflammatory cytokines such as CRP, TNF-, IL-6 and IL-8 [7]. This study was aimed to determine the potential of tender coconut water in inhibiting inflammation due to cigarette smoke exposure.

2 Method

This was an experimental study in a posttest only control group design, twenty male Wistar rats kept in Inter-University Research (PAU) Nutrition Laboratory of PSPG Faculty of Medicine, Gadjah Mada University, Yogyakarta were randomly divided into 4 groups (n=6 rats). Group division and treatment can be seen in Table 1

Table 1 Group division and treatment

Group	Treatment
Negative control (K1)	standard diet <i>et libitum</i> + aqua (morning-afternoon) for 14 days
Positive control (K2)	standard diet <i>et libitum</i> + aqua + cigarette smoke exposure (3 cigarettes/day (morning-afternoon) for 14 days
Treated group 1 (K3)	standard diet <i>et libitum</i> + aqua + cigarette smoke exposure (3 cigarettes/day + vitamin E at the dose of 1.8 mg/200gBW/day (morning-afternoon) for 14 days
Treated group 2 (K4)	Cigarette smoke (3 cigarettes/day + tender coconut water at the dose of 8mL/200gBW/day (morning-afternoon) for 14 days.

On day 15, blood samples were collected for the evaluation of levels of CRP, IL-6, and TNF- α using the ELISA (enzyme-linked immunosorbent assay) method.

2.1 Cigarette Smoke Exposure

The rats were transferred to the smoking chamber (special cage) according to their group when they were exposed to cigarette smoke. The cage was a smoking

box with a barrier to separate the experimental animals from the burning ends of the cigarettes. Cigarette smoke was exhaled repeatedly with the help of an injection tube until the cigarette burns out. The number of cigarettes used in this study was 3 cigarettes/day and administered for 14 days.

2.2 Blood Sampling

The equipment used was sterile micro-hematocrite tubes, blood collection bottles and sterile cotton. Blood sample was taken by inserting a micro-hematocrite tube into the ophthalmic vein in the corner of the rat's eyeball periorbital and then slowly rotated until the blood came out. The blood was accumulated in 2cc Eppendorf. The micro-hematocrite tube was removed after the required blood obtained was sufficient. The blood in the corner of the rat's eyeball was cleaned with sterile cotton.

2.3 Measurement of CRP, IL-6 and TNF- α

The measurement the levels of CRP, IL-6 and TNF α were assessed using the ELISA (*Enzyme-Linked Immunosorbent Assay*) kit.

2.4 Determination of Tender Coconut Water Dosage

The Dosage of tender coconut water was determined based on that of proposed by Zulaikhah et al. ie (4 mL/100gBW).

2.5 Research Place

Treatment of rats and measurement of CRP, IL-6 and TNF α in Inter-University Research (PAU) Nutrition Laboratory of PSPG Faculty of Medicine, Gadjah Mada University, Yogyakarta.

2.6 Data Analysis

The data were tested for normality with Shapiro-Wilk and resulted in a normal and non-homogeneous distribution. Then the data were analyzed using one way ANOVA test followed by the post hoc Tamhane test to compare between treatment groups. The TNF- α level data had a normal and homogeneous distribution, then the data were analyzed by one way ANOVA test followed by Post Hoc LSD test to compare between treatment groups. The decision to accept or reject the hypothesis was based on α 5%.

3 Findings

The effect of tender coconut water on CRP, IL-6 and TNF is shown in Table 2.



Table 2: The effect of tender coconut water on the level of CRP, IL-6 and TNF in the four groups

VARIABLE	GROUP				p-value
	K1	K2	K3	K4	
	Mean ±SD	Mean ±SD	Mean ±SD	Mean ±SD	
CRP Level (ng/mL)	3,2±0,28	17,2±0,86	5,4±0,39	6,7±0,33	>0,05*
Shapiro wilk					<0.05**
Levene test	0,161	0,091	0,988	0,1222	0,0001***
One way ANOVA					
IL-6 Level (pg/mL)	43,0±6,09	87,8±3,87	53,9±1,87	62,8±3,82	>0,05*
Shapiro wilk					<0.05**
Levene test	0,111	0,996	0,616	0,961	0,0001***
One way ANOVA					
TNF- α level (pg/mL)	6,06±0,30	17,9±0,26	7,3±0,39	8,8±0,43	>0,05*
Shapiro wilk					>0.05**
Levene test	0,737	0,858	0,933	0,825	0,0001***
One way ANOVA					

Significant *>0,05
 **>0,05
 ***<0,05

Mean CRP, IL-6 and TNF- α and Post hoc Tamhane and LSD are presented in Figure 1




Fig. 1 Differences in mean CRP levels between groups

**** Significant p<0,0001
 ** Significant p<0,05

Table 1 and Figure 1 show that the highest and the lowest mean levels of CRP were found in the K2 (7.2±0.86 ng/mL) and K1 (3.2±0.28 ng/mL) group, respectively. K3 and K4 (53.9±1.87 pg/mL and 62.8±3.82 pg/mL) had lower mean levels of IL-6 than K2 but higher than K1. The difference in the mean level of IL-6 between K2 and K3 was 11.8 ng/m; between K2 and K4 was 10.5 ng/mL. The results of the ANOVA showed that the administration of tender

coconut water at the dose of 8mL/200gBW/day for 14 days reduced IL-6 levels ($p < 0.05$). The mean IL-6 levels in (K3) was lower compared to that of tender coconut water (K4). The results of the analysis using the Post hoc Tamhane test showed that there was a statistically significant difference in mean IL-6 level between the two groups.




Fig. 2 Differences in mean IL-6 levels between groups

**** Significant ($p < 0.0001$)
 ** Significant ($p < 0.05$)

Table 1 and Figure 2 show that K2 and K1 group had the highest (87.8 ± 3.8 pg/mL) and the lowest (43.0 ± 6.09 pg/mL) mean IL-6 level, respectively. K3 and K4 had lower mean IL-6 level (53.9 ± 1.87 pg/mL and 62.8 ± 3.82 pg/mL) than K2 but higher than K1. The difference in mean level IL-6 levels between K2 and K3 was 33.9 pg/mL, K2 and K4 was 25.0 pg/mL.

The results of the analysis with ANOVA showed that the administration of tender coconut water at the dose of 8mL/200gBW/day for 14 days reduced IL-6 levels ($p < 0.05$). The mean IL-6 levels in K3 was lower compared to K4. The results of the analysis using the Post hoc Tamhane test showed that there was a statistically significant difference in mean IL-6 levels between the two groups.




Fig. 3 Differences in mean TNF-α levels between groups

**** Significant ($p < 0.0001$)
 ** Significant ($p < 0.05$)

Table 1 and Figure 3 show that the highest and the lowest mean levels of TNF-α was found in K2 group (17.9 ± 0.26 pg/mL) and K1 (6.06 ± 0.30 pg/mL), group respectively. K3 (7.3 ± 0.39 pg/mL) and K4 (8.8 ± 0.43 pg/mL) had lower mean levels than K2 but higher than K1. The difference in mean level of TNF-α between K2 and K3 was 10.6 pg/mL; K2 and K4 was 9.1 pg/mL.

The results of the analysis with ANOVA showed that the administration of tender coconut water at the dose of 8mL/200gBW/day for 14 days reduced IL-6 levels ($p < 0.05$). The mean level of IL-6 in K3 group was lower compared to K4. The analysis using the Post hoc Tamhane test showed that there was a statistically significant difference in mean levels of IL-6 between the two groups.

4 Discussion

The results of this study showed that mean level of CRP, IL-6 and TNF-α in the group exposed to cigarette smoke were higher than that of group without the exposure of cigarette smoke. This indicates that tender coconut water has a potential in inhibiting inflammation. These findings are consistent with that of the previous studies showing that smoking increased levels of CRP and IL-6, the presence of an inflammatory reaction increases levels of CRP, white blood cells, albumin, interleukin 6 (IL-6) and tumor necrosis factor (TNFα) in the blood [8]. Long-term exposure to cigarette smoke can cause airway inflammation characterized by increased neutrophils, macrophages, T lymphocyte counts and cytokines such as tumor necrosis factor-alpha (TNF-α), interleukin (IL)-6 and IL-8 [9]. The production of cytokines are mediated, especially by free radicals derived from oxygen. These free radical mediators help the formation of cytokines by modulating nuclear factor (NFκB). Cigarette smoke contains free radical that causes oxidative stress. Cigarette smoke-induced oxidative stress causes a series of cellular and molecular reactions, activates cascades and transcription factors, releases inflammatory mediators, initiates inflammation, cell injury and apoptosis [10].

One puff of cigarette smoke contains numerous free radicals which plays a role in the pathogenesis. The major free radicals through cigarette smoke are superoxide (O_2^-), nitric oxide (NO), which combine to form peroxynitrite [11]. Due to the accumulation of free radicals, tissues are indented to deplete antioxidants such as ascorbic acid and protein

sulfhydryl groups leading to the oxidation of DNA, lipids, and proteins. This ROS accumulation in tissues might be due to an increase in oxidant generation, a decrease in antioxidant defence agents, or failure to repair oxidative damage. The imbalance between oxidants and antioxidants causes oxidative stress initiating the inflammatory response [4]. The response begins with an increased macrophages, neutrophils and T lymphocytes. This triggers the release of pro-inflammatory cytokines such as tumor necrosis factor- α (TNF- α), interleukin (IL)-6 and IL-8 [7]. Several studies have shown increased levels of CRP, TNF-, IL-6 and IL-8 in COPD patients with inflammation due to exposure to cigarette smoke. Levels of C-reactive protein (CRP), TNF-, IL-6 and IL-8 in smoking COPD patients were shown to be higher than that of non-smoking. The exposure of cigarette smoke increases levels of pro-inflammatory cytokines [5], [12]. CRP, IL-6 and TNF- α levels are pro-inflammatory cytokines. Increased CRP, IL6 and TNF- α level indicate high levels of ROS and decreased nitrate oxidation. High levels of IL-6 stimulates the liver to synthesize and secrete markers of systemic inflammation including CRP [13].

The most dangerous content of cigarette smoke causing oxidative stress and inflammation is nicotine, carbon monoxide, metals (arsenic, nickel, chromium and lead) and particulate matter and acrolein. Nicotine in cigarettes can cause an increase in ROS and a decrease in the antioxidant capacity of enzymes, as well as induce the formation of active compounds disrupting blood flow by inducing inflammation in blood vessels. Carbon monoxide in cigarettes can cause hypoxia and an increase in xanthine oxidase and nicotinamide adenine dinucleotide phosphate oxidase (NADPH Oxidases). Acrolin in cigarettes can cause a decrease in antioxidant enzymes, decrease in the immune response and an increase in inflammation markers [14].

Without an excessive amount of antioxidant reserves, the body requires higher exogenous antioxidants to eliminate the effects of free radicals exposure [15]. There are different enzymatic mechanisms to minimize radical damage and to protect against excessive production of free radicals. Antioxidants play an important role in this defence mechanism [16]. In normal and healthy subjects, there is a balance between the formation of reactive oxygen species free radicals and endogenous antioxidant defense mechanisms [17], if ROS amount more than antioxidants, then the excessive amount of ROS will disturb lipid, protein, and DNA components, called oxidative stress [15]. At present, the impact of oxidative stress and its related factors has become an important issue for human health. The World Health Organization (WHO) estimates that about 80% of the world's population uses traditional medicine for

primary health care. In recent years, more interest has been focused on experimental research to verify natural antioxidants and anti-inflammatory drugs from natural resource products [18]. Foods containing natural antioxidants can be used as a strategy to reduce morbidity and mortality, especially due to oxidative stress [15]. Natural source of antioxidants can be used as preventive medicine. Recent studies have shown that there is an inverse relationship between consumption of antioxidant-rich foods and the prevalence of human disease. Consuming antioxidants from natural plant sources as daily diet can offer solution to overcome human health problems [17], [19].

Tender coconut water contains various nutrients such as L-arginine, Vitamin C, amino acids and minerals such as magnesium, potassium, calcium, selenium, methionine, zinc, iodine, manganese, and cuprum [15], [20]. The results of this present study showed that the administration of tender coconut has a potential to inhibit inflammation due to cigarette smoke indicated by lower levels of CRP, IL-6 and TNF in the K4 group when compared to the group exposed to cigarette smoke. The results of this study supports that of a study conducted by Zulaikhah et al showing that administration of tender coconut water at the dose of 8mL/200gr BW for 4 weeks prevents inflammation characterized by a decrease in TNF α , IL-1 and IL-6 in induced by streptozotocin (STZ) and nicotinamid (NA) [8].

Vitamin C in tender coconut water serving as an antioxidant, is a micronutrient that plays an important role in the immune system with pleiotropic functions due to its ability to donate electrons, biosynthesis of enzymes and function of gene regulation. Vitamin C also playing a role in cellular immunity, increasing epithelial barrier function, is needed in the apoptosis and has the ability to cleanse necrotic tissue (NETosis) to reduce the inflammatory process [21]. A study conducted by Dewi et al. proved that Vitamin C reduce pro-inflammatory cytokines characterized by differences in IL-6 and CRP levels between the group a with vitamin C at a dose of 12 g/50 mL water every 12 hours for 7 days and the placebo group showing that mean levels of IL-6 and CRP in the group given vitamin C was lower compared to placebo group [22].

Vitamin C has been shown to decrease nicotine toxicity by donating hydrogen atom to nicotine metabolite indicated by the difference in CE expression between group supplemented with Vitamin C at the doses of 200 mg/day and 1000 mg/day compared with placebo. The decrease in CE excretion positively correlated with the increased dose of vitamin C supplementation [23].

Vitamin C has been shown to decrease inflammation and oxidative stress in patients with metabolic syndrome indicated by decreased CRP dan IL-6 level [24]. Animal studies have shown that vitamin C

deficiency exposed to cigarette smoke indicates the release of Troponin T and I in serum, oxidative stress, apoptosis, thrombosis and myocardial deposition. There was a significant difference in plasma vitamin C levels between smokers and non-smoker group [25].

Vitamin C is a micronutrient that plays an important role in the immune system with pleiotropic functions due to its ability to donate electrons, biosynthesis of enzymes and function of gene regulation. Vitamin C also functions in cellular immunity, increases epithelial barrier function, is needed in the process of apoptosis and has the ability to cleanse necrotic tissue (NETosis) to reduce the inflammatory process [23]. Research on experimental showed that vitamin C deficiency exposed to cigarette smoke showed the release of Troponin T and I in serum, oxidative stress, apoptosis, thrombosis and myocardial deposition. There was a significant difference in plasma vitamin C levels in smokers compared to the non-smoker group [25].

CRP levels are inversely correlated with vitamin C levels in the body, meaning that the higher the intake of vitamin C, the lower the levels of CRP will be and the lower the intake of vitamin C, the higher the levels of CRP will be [26]. Vitamin C acts as an antioxidant and immunomodulator in reducing the severity of COVID-19 infection [22]. The reduction of free radicals can prevent oxidative stress so that it can inhibit the occurrence of inflammatory responses in the body. Vitamin C can also reduce the inflammatory response directly by suppressing the production of IL-6, CRP, and TNF- α [13], [22].

L-arginine in tender coconut water is a non-essential amino acid, in nitric oxide synthase (NO Synthase) this compound is a substrate to produce citrulline and NO. NO can inhibit xanthine oxidase (XO) leading to increased SOD levels, total thiol levels (T-SH), vitamin C, total antioxidants (TAC) [20]. The results of previous studies showed that L-arginine inhibited the secretion of IL-1 β , IL-6, and TNF- by macrophages and other inflammatory infiltrating cells in the male dystrophin-deficient (MDX) diaphragm muscle. L-arginine inhibits the inflammatory cascade so as to reduce the number and activity of nuclear factor-kB (NF-kB) and metalloproteinase (MMP)-2 and MMP-9 activities in male dystrophin-deficient (MDX) mice [27].

L-arginine in tender coconut water is antioxidant and can reduce free radicals characterized by increased levels of SOD, catalase and GPx [20]. L-arginine will bind to free radicals directly or damage free radical bonds [28]. L-arginine compounds can inhibit inflammation as seen from the decrease in levels of pro-inflammatory cytokines such as CRP, TNF-, IL-6 and IL-8 [8].

5 Conclusion

Cigarette smoke can cause stress oxidative that can induce inflammatory responses [12], [14]. Tender coconut water consist of L-arginin, Vitamin C, and amino acids that can inhibits inflammation by decreasing levels of pro-inflammatory cytokines such as CRP, IL-6, IL-8, and TNF α [8], [19].

There are several studies that concern on the effect of tender coconut water. The oral administration of tender coconut water could lower TC, LDL and TG levels and increase HDL serum levels in male rats fed high fat diet [15]. Other research shown consumption of tender coconut water for 30 days increases antioxidant enzymatic SOD, CAT, GPx, and decreases lipid peroxidation in mercury exposure workers [20]. The administration of tender coconut water can decrease the levels of TNF- α , IL-1 and IL-6 in Streptozotocin (STZ) and Nicotinamide (NA) induced diabetic rats [8]. Coconut water can effectively reduce the level of heavy metal, especially Cd and Hg, in the blood of *Rattus norvegicus* [28].

This present study showed that the administration of tender coconut water has a potential effect to inhibit inflammation due to cigarette smoke. This study demonstrated that the administration of tender coconut water was able to inhibits inflammation due to cigarette smoke, which was characterized by decreased levels of CRP, IL-6 and TNF- α in the K4 group when compared to K2 in mice exposed to cigarette smoke.

The limitation in this study is that the study only uses tender coconut water with one dose and one type of tender coconut water. Despite its limitation, this study has strength including its study aim that find out the effect of tender coconut water in inhibits inflammation caused specifically by cigarette smoke, which is rarely studied. We suggest that researchers need to investigate the effect of tender coconut water in inhibits inflammation due to cigarette smoke using various dosage and different types of tender coconut water for further research.

References

- [1] H. HOLIPAH, H. W. SULISTOMO, AND A. MAHARANI, Tobacco smoking and risk of all-cause mortality in Indonesia, *PLoS One*, vol. 15, no. 12 December, pp. 1–12, 2020, <https://doi.org/10.1371/journal.pone.0242558>
- [2] CENTRES FOR DISEASE CONTROL AND PREVENTION, Smoking and Tobacco Use : Data and Statistic, 2021. https://www.cdc.gov/tobacco/data_statistics/

- patients, 2020. *IP Indian J. Immunol. Respir. Med.*, vol. 5, no. 3, pp. 173–180. 10.18231/j.ijirm.2020.053
- [3] WORLD HEALTH ORGANIZATION, Country Profile Indonesia WHO Report on the Global Tobacco Epidemic 2019, *World Health Organization.*, pp. 1–10. 2019. https://www.who.int/tobacco/surveillance/policy/country_profile/idn.pdf
- [4] A. STRZELAK, A. RATAJCZAK, A. ADAMIEC, AND W. FELESZKO, Tobacco smoke induces and alters immune responses in the lung triggering inflammation, allergy, asthma and other lung diseases: A mechanistic review. *Int. J. Environ. Res. Public Health.*, vol. 15, no. 5. 2018. 10.3390/ijerph15051033
- [5] R. V. SURYADINATA, Pengaruh Radikal Bebas Terhadap Proses Inflamasi pada Penyakit Paru Obstruktif Kronis (PPOK), *Amerta Nutr.*, vol. 2, no. 4, p. 317, 2018. 10.20473/amnt.v2i4.2018.317-324
- [6] Z. LEI, H. GUO, S. ZOU, J. JIANG, Y. KUI, AND J. SONG, “Long non-coding RNA maternally expressed gene regulates cigarette smoke extract induced lung inflammation and human bronchial epithelial apoptosis via miR-149-3p,” *Exp. Ther. Med.*, vol. 21, no. 1, pp. 1–9, 2020. 10.3892/etm.2020.9492
- [7] E. YI *et al.*, “Long noncoding RNA IL6-AS1 is highly expressed in chronic obstructive pulmonary disease and is associated with interleukin 6 by targeting miR-149-5p and early B-cell factor 1, 2021. *Clin. Transl. Med.*, vol. 11, no. 7. 10.1002/ctm2.479
- [8] ZULAIKHAH, S.T., J. WAHYUWIBOWO, M. N. SUHARTO, B. H. ENGGARTIASTO, M. I. R. ORTANTO, AND A. A. PRATAMA, Effect of tender coconut water (TCW) on TNF- α , IL-1 and IL-6 in streptozotocin (STZ) and nicotinamid (NA) induced diabetic rats, *Pharmacogn. J.*, vol. 13, no. 2, pp. 500–505. 2021. 10.5530/pj.2021.13.63
- [9] N. A. KHAN *et al.*, “Systemic Biomarkers of Inflammation, Oxidative Stress and Tissue Injury and Repair among Waterpipe, Cigarette and dual Tobacco Smokers,” *Tob. Control J.*, Vol 29, pp. 102-109. 2020. 10.1136/tobaccocontrol-2019-054958
- [10] H. LAWRENCE, A. HUNTER, R. MURRAY, W. S. LIM, AND T. MCKEEVER, “Cigarette Smoking And The Occurrence Of Influenza – Systematic Review,” *J. Infect.*, vol. 79, no. 5, pp. 401–406, 2019. 10.1016/j.jinf.2019.08.014
- [11] S. ASEERVATHAM, S. CHOI, J. KRIHNNAN, AND R. K. Cigarette Smoke and Related Risk Factors in Neurological Disorders: An update., 2017. *Biomed Pharmacother.*, vol. 85, pp. 79–86. <https://doi.org/10.1016/j.biopha.2016.11.118>
- [12] M. ASIMUDDIN, G. VIJAYLAKSHMI, M. SOHEB ANSARI, AND K. JAMIL, Biomarkers of inflammation and oxidative stress in smokers and severe chronic obstructive pulmonary disease
- [13] Q. T. TRAN, S. ARIMILLI, E. SCOTT, P. CHEN, AND G. L. PRASAD, “Differences in Biomarkers of Inflammation and Immune Responses in Chronic Smokers and Moist Snuff Users,” *Cytokine*, vol. 137, no. June 2020, p. 155299, 2021. <https://doi.org/10.1016/j.cyto.2020.155299>
- [14] B. TAATI, H. ARAZI, AND K. SUZUKI, “Oxidative stress and inflammation induced by waterpipe tobacco smoking despite possible protective effects of exercise training: A review of the literature, 2020. *Antioxidants*, vol. 9, no. 9, pp. 1–13. 10.3390/antiox9090777
- [15] ZULAIKHAH, S.T., D. PERTIWI, S. A. BAGUS, S. NURI, E. M. BRILLIAN JELITA, AND N. S. ALFIZA, Effect of tender coconut water on blood lipid levels in high fat diet fed male rats, 2017. *J. Krishna Inst. Med. Sci. Univ.*, vol. 6, no. 2, pp. 63–68.
- [16] M. SHARIFI-RAD *et al.*, Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases, 2020. *Front. Physiol.*, vol. 11, no. July, pp. 1–21. 10.3389/fphys.2020.00694
- [17] P. ARULSELVAN *et al.*, Role of Antioxidants and Natural Products in Inflammation, 2016 *Oxid. Med. Cell. Longev.*, vol. 2016. 10.1155/2016/5276130
- [18] KEMENTRIAN KESEHATAN RI, Hasil Riset Kesehatan Dasar Tahun 2018, 2018 *Kementrian Kesehat. RI*, vol. 53, no. 9, pp. 1689–1699.
- [19] ZULAIKHAH, S.T., Health Benefits of Tender Coconut Water (TCW), 2019. *Int. J. Pharm. Sci. Res.*, vol. 10, no. 2, pp. 474–480. 10.13040/IJPSR.0975-8232.10(2).474-80
- [20] ZULAIKHAH, S.T., AND A. SUWONDO, Effects of Tender Coconut Water on Antioxidant Enzymatic Superoxida Dismutase (SOD), CATALASE (CAT), Glutathione Peroxidase (GPx) and Lipid Peroxidation In Mercury Exposure Workers, 2017 *Int. J. Sci. Res.*, vol. 4, no. 12, pp. 517–524. 10.21275/v4i12.nov151788
- [21] A. C. CARR AND S. MAGGINI, “Vitamin C and immune function, 2017. *Nutrients*, vol. 9, no. 11, pp. 1–25. <http://10.3390/nu9111211>
- [22] A. M. C. DEWI, E. M. DAGRADI, AND P. WIBOWO, The Effect of High Dose Vitamin C (ascorbic acid) on Pro inflammatory Cytokines in COVID-19, 2021. *Med. Heal. Sci. J.*, vol. 5, no. 1, pp. 46–50. 10.33086/mhsj.v5i1.1884
- [23] P. HOLFORD *et al.*, “Vitamin C—An adjunctive therapy for respiratory infection, sepsis and COVID-19,” *Nutrients*, vol. 12, no. 12, pp. 1–17, 2020. 10.3390/nu12123760
- [24] S. K. WONG, K. Y. CHIN, AND S. IMANIRWANA, Vitamin C: A review on its role in the management of metabolic syndrome, 2020 *Int. J.*

- Med. Sci.*, vol. 17, no. 11, pp. 1625–1638. 10.7150/ijms.47103
- [25] G. XIA, D. FAN, Y. HE, Y. ZHU, AND Q. ZHENG, “High-Dose Intravenous Vitamin C Attenuates Hyperinflammation In Severe Coronavirus Disease 2019,” *Nutrition*, vol. 91–92, p. 111405, 2021. <https://doi.org/10.1016/j.nut.2021.111405>
- [26] I. JIALAL AND U. SINGH, Is vitamin C an antiinflammatory agent?. 2016 *Am. J. Clin. Nutr.*, vol. 83, no. 3, pp. 525–526. 10.1093/ajcn.83.3.525
- [27] K. HNIA *et al.*, L-arginine decreases inflammation and modulates the nuclear factor- κ B/matrix metalloproteinase cascade in Mdx muscle fibers, 2018 *Am. J. Pathol.*, vol. 172, no. 6, pp. 1509–1519. 10.2353/ajpath.2008.071009
- [28] Y. RACHMAWATI *et al.*, Effectiveness of Green Coconut (Cocos nucifera L.) Water against Heavy Metal Levels in the Blood of Rattus norvegicus. 2020. pp. 113–118., <http://10.5220/0008907201130118>.

15. SITI THOMAS_Tender Coconut Water can Inhibits Inflammation due to Cigarette Smoke

ORIGINALITY REPORT

17%

SIMILARITY INDEX

12%

INTERNET SOURCES

12%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	digitalcommons.unmc.edu Internet Source	1%
2	Pedro Lourenco Katayama, Isabela de Paula Leirao, Alexandre Kanashiro, Joao Paulo Mesquita Luiz et al. "The Carotid Body Detects Circulating Tumor Necrosis Factor-Alpha to Activate a Sympathetic Anti-Inflammatory Reflex", Cold Spring Harbor Laboratory, 2021 Publication	1%
3	gssrr.org Internet Source	<1%
4	pesquisa.bvsalud.org Internet Source	<1%
5	www.ijbcp.com Internet Source	<1%
6	D. S. Aniesrani Delfiya, K. Thangavel. "Effect of Ohmic Heating on Polyphenol Oxidase Activity, Electrical and Physicochemical	<1%

Properties of Fresh Tender Coconut Water",
International Journal of Food Engineering,
2016

Publication

7

Titik Sumarawati, - Israhnanto, Dina
Fatmawati. "Anticancer Mechanism of Melia
azedarach, Doxorubicin and Cyclophosphamide
Combination against Breast Cancer in Mice",
Bangladesh Journal of Medical Science, 2017

Publication

<1 %

8

escholarship.org

Internet Source

<1 %

9

www.frontiersin.org

Internet Source

<1 %

10

www.scitepress.org

Internet Source

<1 %

11

Altinors, D.D.. "Smoking Associated With
Damage to the Lipid Layer of the Ocular
Surface", American Journal of Ophthalmology,
200606

Publication

<1 %

12

Dipak P. Pandya. "Oxidant injury and
antioxidant prevention: Role of dietary
antioxidants, minerals, and drugs in the
management of coronary heart disease [Part
II]", Comprehensive Therapy, 2002

Publication

<1 %

13

Ruoming Tan, Xiaogang Xiang, Wei Chen, Zhitao Yang, Weiguo Hu, Hongping Qu, Jialin Liu. "Efficacy of diammonium glycyrrhizinate combined with vitamin C for treating hospitalized COVID-19 patients: a retrospective, observational study", QJM: An International Journal of Medicine, 2021

Publication

<1 %

14

bioone.org

Internet Source

<1 %

15

www.thieme-connect.com

Internet Source

<1 %

16

Sinha, M.. "A 43kD protein from the herb, *Cajanus indicus* L., protects against fluoride induced oxidative stress in mice erythrocytes", Pathophysiology, 20070531

Publication

<1 %

17

www.ahvma.org

Internet Source

<1 %

18

Dewa Ketut Meles, Imam Mustofa, Wurlina Wurlina, Suherni Susilowati et al. "The Restorative Effect of Red Guava (*Psidium guajava* L.) Fruit Extract on Pulmonary Tissue of Rats (*Rattus norvegicus*) Exposed to Cigarette Smoke", Veterinary Medicine International, 2021

Publication

<1 %

19	Handa, James T.. "How does the macula protect itself from oxidative stress?", <i>Molecular Aspects of Medicine</i> , 2012. Publication	<1 %
20	www.oncotarget.com Internet Source	<1 %
21	Suparmi Suparmi, Diana Widiastuti, Sebastiaan Wesseling, Ivonne M.C.M. Rietjens. "Natural occurrence of genotoxic and carcinogenic alkenylbenzenes in Indonesian jamu and evaluation of consumer risks", <i>Food and Chemical Toxicology</i> , 2018 Publication	<1 %
22	Submitted to University College London Student Paper	<1 %
23	lib.fkik.untad.ac.id:80 Internet Source	<1 %
24	pure-oai.bham.ac.uk Internet Source	<1 %
25	www.dovepress.com Internet Source	<1 %
26	Bidisha Rajkhowa, Sidharth Mehan, Pranshul Sethi, Sonalika Bhalla et al. "Solanesol Mediated SIRT-1 Activation Prevents Neurobehavioral and Neurochemical Defects in Ouabain-Induced Experimental Model of	<1 %

Bipolar Disorder in Rats", Research Square Platform LLC, 2021

Publication

27

Nicole Greisman, Bianca Wollenhaupt-Aguiar, Taiane de Azevedo Cardoso, Karen Jansen et al. "CCL11 levels in drug-naive bipolar patients: The role of sex and smoking status", Journal of Affective Disorders Reports, 2020

Publication

<1 %

28

Submitted to University of Birmingham

Student Paper

<1 %

29

dentistry.tbzmed.ac.ir

Internet Source

<1 %

30

ijms.sums.ac.ir

Internet Source

<1 %

31

www.banglajol.info

Internet Source

<1 %

32

www.spandidos-publications.com

Internet Source

<1 %

33

"Environmental Contaminants and Neurological Disorders", Springer Science and Business Media LLC, 2021

Publication

<1 %

34

A. S. Mohan Kumar, V. Geetha, G. Suresh Kumar. "Impact of phenolic concentrate from tender coconut water and coconut testa on

<1 %

high fat cholesterol fed C57BL6 mice", Journal of Food Science and Technology, 2021

Publication

35

Daniela Miglietta, Clara De Palma, Clara Sciorati, Barbara Vergani et al. "Naproxcinod shows significant advantages over naproxen in the mdx model of Duchenne Muscular Dystrophy", Orphanet Journal of Rare Diseases, 2015

Publication

<1 %

36

Sandhya Vijayam Gopalakrishnan Nair, Thankappan Rajamohan. "The Role of Coconut Water on Nicotine-Induced Reproductive Dysfunction in Experimental Male Rat Model", Food and Nutrition Sciences, 2014

Publication

<1 %

37

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

38

www.hindawi.com

Internet Source

<1 %

39

Akio Kawamura, Masao Saitoh, Motoki Yonekawa, Takashi Horie, Hiroko Ohizumi, Tohru Tamaki, Kazutaka Kukita, Jun - ichi Meguro. "New Technique of Leukocytapheresis by the Use of Nonwoven Polyester Fiber Filter for Inflammatory Bowel Disease", Therapeutic Apheresis, 2002

<1 %

40

Philip Thomas James, Zakari Ali, Andrew E Armitage, Ana Bonell et al. "Could nutrition modulate COVID-19 susceptibility and severity of disease? A systematic review", Cold Spring Harbor Laboratory, 2020

Publication

41

S. Teramoto, Y. Uejima, K. Teramoto, Y. Ouchi, Y. Fukuchit. "Effect of age on alteration of glutathione metabolism following chronic cigarette smoke inhalation in mice", Lung, 1996

Publication

42

V L Kinnula. "Focus on antioxidant enzymes and antioxidant strategies in smoking related airway diseases", Thorax, 2005

Publication

43

actanaturae.ru

Internet Source

44

bsdwebstorage.blob.core.windows.net

Internet Source

45

docsdrive.com

Internet Source

46

dokumen.pub

Internet Source

47

dranthonylim.wordpress.com

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

48

dro.deakin.edu.au

Internet Source

<1 %

49

dx.doi.org

Internet Source

<1 %

50

ijphs.iaescore.com

Internet Source

<1 %

51

journal2.unusa.ac.id

Internet Source

<1 %

52

juniperpublishers.com

Internet Source

<1 %

53

jurnal.unissula.ac.id

Internet Source

<1 %

54

peerj.com

Internet Source

<1 %

55

Bernard Fischer, Judith Voynow, Andrew Ghio.
"COPD: balancing oxidants and antioxidants",
International Journal of Chronic Obstructive
Pulmonary Disease, 2015

Publication

<1 %

56

S J Kloprogge. "Expression of TSH-R in normal
human extraocular muscles", British Journal
of Ophthalmology, 1/1/2006

Publication

<1 %

57

Yea Seong Ryu, Kyoung Ah Kang, Mei Jing Piao, Mee Jung Ahn et al. "Particulate matter induces inflammatory cytokine production via activation of NFκB by TLR5-NOX4-ROS signaling in human skin keratinocyte and mouse skin", Redox Biology, 2019

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On

15. SITI THOMAS_Tender Coconut Water can Inhibits Inflammation due to Cigarette Smoke

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
