

DAFTAR PUSTAKA

- Alvarez-Llamas G, Szalowska E, Vries MP, Weening D, Landman K, Hoek A, Wolffenduttel BH, Roelofsen H, Vonk RJ: Characterization of the human visceral adipose tissue secretome. *Mol Cell Proteomics*, 2007;6:589–600. doi: 10.1074/mcp.M600265-MCP200.
- Bruno MB, Antonio CAN, Francelius DO: Mesenchymal Stem Cells - Emphasis in Adipose Tissue, *Brazilian Archives of Biology and Technology*, 2013: 607-617.
- Considine RV: Regulation of leptin production. *Rev.endocrine*. 2001. 357-363.
- Dexheimer V, Frank S, Richter W: Proliferation as a requirement for in vitro chondrogenesis of human mesenchymal stem cells. *Stem Cells Dev*, 2012. 21:2160–2169.
- Dhanasekaran M, Indumathi S, Lissa RP, Harikrishnan R, Rajkumar JS, Sudarsanam D: A comprehensive study on optimization of proliferation and differentiation potency of bone marrow derived mesenchymal stem cells under prolonged culture condition. *Cytotechnology*. 2013;65:187–197.
- Dhanasekaran M, Indumathi S, Rashmi M, Rajkumar JS, Sudarsanam D: Unravelling the retention of proliferation and differentiation potency in extensive culture of human subcutaneous fat-derived mesenchymal stem cells in different media. *Cell Prolif* 2012, 45:516–526.
- Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, Deans R, Keating A, Prockop D, Horwitz E: Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy*, 2006, 8:315–317.
- Edwin C, Mariman M, Wang P: Adipocyte extracellular matrix composition, dynamics and role in obesity. *Cell Mol Life Sci*. 2010, (8): 1277–1292
- Fernandez VB, Romaniuk MA, Choi H, Labovsky V, Otaegui JC: Mesenchymal stem cells and their use in therapy: what has been achieved?. *Differentiation*, 2013: 1-10
- Gomillion CT, Burg KJ: Stem cells and adipose tissue engineering. *Biomaterials*, 2006; 27(36):6052-63.
- Hagmann S, Moradi B, Frank S, Dreher T, Kammerer PW: Different culture media affect growth characteristics, surface marker distribution and chondrogenic differentiation of human bone marrow-derived mesenchymal stromal cells. *BMC Musculoskeletal Disorders*, 2013;(14) 1471-2474.

Houseknecht KL, Cole BM, Steele PJ: Peroxisome proliferatoractivated receptor gamma (PPAR γ) and its ligands: a review. *Domest.Anim.Endocrinol.* 2002; 22, 1-23.

Huss FRM, Kratz G. Adipose tissue processed for lipoinjection shows increased cellular survival *in vitro* when tissue engineering principles are applied. *Scand.J.Plast.Reconstr.Surg.* 2002; 36, 166-171.

Ikawati, Z. 2006. *Pengantar Farmakologi Molekuler*. Yogyakarta: Gadjah Mada University Press

Kim YC, Gomez FE, Fox BG and Ntambi JM: Differential regulation of the stearoyl-CoA desaturase genes by thiazolidinediones in 3T3-L1 adipocytes. *J.Lipid Res.* 2000; 41, 1310-1316

Krahenbuhl SM, Grognuz A, Michetti M, Raffoul W, Applegate LA: Enhancement of Human Adipose-Derived Stem Cell Expansion and Stability for Clinical use. *Int J Stem Cell Res.* 2015.

Niemela S, Miettinen S, Konttinen S, Waris T, Kellomaki M, Ashammakhi N, Ylikomi T: Fat Tissue: Views on reconstruction and exploitation. *J Craniofac Surg*, 2007; 18(2): 325-335.

Ntambi JM, Young-Cheul K: Adipocyte differentiation and gene expression. *J Nutr*, 2005; 130: 3122S-3126S

Reusch JE, Colton LA, Klemm DJ: CREB activation induces adipogenesis in 3T3-L1 cells. *Mol Cell Biol*, 2000; 20(3): 1008-20.

Rosen ED, Spiegelman BM: Molecular regulation of adipogenesis. *Annu.Rev.Cell Dev.Biol.* 2000; 16, 145-171.

Rosen ED, MacDougald OA: Adipocyte differentiation from the inside out. *Nat Rev Mol Cell Biol*, 2006; 7:885-896

Sillat T, Saat R, Pollanen R, Hukkanen M, Takagi M, Kottinen YT: Basement membrane collagen type IV expression by human mesenchymal stem cells during adipogenic differentiation. *J Cell Mol Med*, 2012 Jul;16(7):1485-95. doi: 10.1111/j.1582-4934.2011.01442.x.

Sorrell JM, Baber MA, Traktuev DO, March KL, Caplan AI: The creation of an *in vitro* adipose tissue that contains a vascular adipocyte complex. *Biomaterials*, 2011; 32(36): 9667-9676.

Staiger H, Haring HU, Loeffler G: Serum-free differentiation of 3T3-L1 preadipocytes is characterized by only transient expression of peroxisome

proliferatoractivated receptor-gamma. *Biochem.Biophys.Res.Commun*, 2002. 296, 125-128.

Stanners CP, Goldberg VP. N.p., 2016. Web www.sigmaaldrich.com. Jan. 2016.

Sylvestar S, Pathalam G, Dorairaj S, Savariar V: Mesenchymal Stem Cells (MSC): Identification, Proliferation, and Differentiation. *PeerJ Pre Prints*, 2013. doi: 10.7287/peerj.preprints.148v1

Tang QQ, Otto TC, Lane MD: Mitotic clonal expansion: A synchronous process required for adipogenesis. *Proc.Natl.Acad.Sci.U.S.A*, 2003. 100, 44-49.

Walker M, Patel K, Stappenbeck T: The stem cell niche. *J Pathol*. 2009; 217:169-80