

DAFTAR PUSTAKA

1. Escamilla RP, Radovan IM, Dewey KG. The association between caesarean delivery and breast-feeding outcomes among Mexican women. *Am J Public Health*. 1996;86(6):832-6.
2. Carvalho FAE, Tenorio SB. Comparative study between doses of intrathecal morphine for analgesia after caesarean. *Rev Bras Anesthesiol*. 2013;63(6):492-8.
3. Dahl JB, Moiniche S. Pre-emptive analgesia. *British Medical Bulletin*. 2004;71:13-27.
4. Weinger M. Dangers of postoperative opioids. *APFS Newsletter*. 2007;21:61-8.
5. Betran AP, Merialdi M, Lauer JA, Bing-Shun W, Thomas J, Van Look P, et al. Rates of caesarean section: analysis of global, regional and national estimates. *Paediatr Perinat Epidemiol*. 2007;21(2):98-113.
6. National Institute for Health Research and Development. Final report of national basic health research 2010. In: Development NifHRa, editor. Jakarta: The Institute; 2010.
7. Vercauteren M. Analgesia after Caesarean section: are neuroaxila techniques outdated? *Jurnalul Roman de Anestezie Terapie Intensiva*. 2009;16(2):129-33.
8. Ismail S, Afshan G, Monem A, Ahmed A. Postoperative Analgesia Following Caesarean Section: Intravenous Patient Controlled Analgesia versus Conventional Continuous Infusion. *Open Journal of Anesthesiology*. 2012;2:120-6.
9. Verstraete S, Velde MVD. Post-caesarean section analgesia. *Acta Anaesth Belg*. 2012;63:147-67.
10. Man A. Management of Postoperative Pain Hongkong: Department of Anesthesia & Intensive Care The Chinese University of Hong Kong Prince of Wales Hospital; 2014 [cited 2014 2014.4.14]. Available from: <http://www.cuhk.edu.hk/med/ans/pain/Pain Nausea and vomiting.doc>.
11. Morrison R, Magaziner J, McLaughlin M, Orosz G, Silberzweig S, Koval K, et al. The impact of postoperative pain on outcomes following hip fracture. *Pain*. 2003;103(3):303-11.
12. Aoki Y, Hatakeyama N, Yamada M, Matsuda N, Yamazaki M. Postoperative pain management by continuous intravenous infusion of fentanyl using the single -use continuous infusion device. *Masui*. 2009;58(11):1424-9.
13. Kim JY, Park SY, Chang HS, Nam SK, Min SK. The efficacy of the time-scheduled decremental continuous infusion of fentanyl for postoperative patient-controlled analgesia after total intravenous anesthesia. *Korean J Anesthesiol*. 2013;65(6):544-51.
14. Coll Am, Ameen JR, Mead D. Postoperative pain assessment tools in day surgery: literature review. *J Adv Nurs*. 2004;46(2):124-33.

15. Desborough JP. The stress response to trauma and surgery. *British Journal of Anaesthesia*. 2000;85:109-17.
16. Salmah S, Choy YC. Comparison of morphine with fentanyl added to intrathecal 0.5% hyperbaric bupivacaine for analgesia after caesarean section *Med J Malaysia* 2009;64(1).
17. IASP. IASP Taxonomy Wasington, D.C. USA: International Association for The Study of PAin; 2012 [cited 2014 2014.4.14]. Available from: <https://http://www.iasp-pain.org/Education/Content.aspx?ItemNumber=1698 - Pain>.
18. Woolf CJ, Salter MW. Neuronal Plasticity: increasing the gain in pain. *Science*. 2000;288(5472):1765-9.
19. McKeen MJ, Quraishi SA. Clinical review of intravenous opioids in acute care. *Journal of Anesthesiology and Clinical Science*. 2013;2(1):1-11.
20. Zichterman A. Opioid Pharmacology and Considerations in Pain Management: U.S. Pharmacist; 2007 [cited 2014 4.19]. Available from: http://www.uspharmacist.com/continuing_education/ceviewtest/lessonid/105473/.
21. Gourlay GK, Kowalski SR, Plummer JL, Cousins MJ, Armstrong PJ. Fentanyl blood concentration-analgesic response relationship in the treatment of postoperative pain. *Anesth Analg*. 1988;67(4):329-37.
22. Viscusi RE, Schetcher LN. Patient -controlled analgesia: Finding a balance between cost and comfort. *AJHP*. 2006;63(8):S1-16.
23. Brill S, Gurman GM, Fisher A. A history of neuroaxial administration of local analgesic and opioids. *Eur J Anaesthesiol*. 2003;20(9):682-9.
24. Bujedo BM. A Clinical Approach to Neuraxial Morphine for the Treatment of Postoperative Pain. *Pain Research and Treatment*. 2012;2012(612145).
25. Rathmell JP, Lair TR, Nauman B. The Role of Intrathecal Drugs in the Treatment of Acute Pain. *Anesth Analg*. 2005;101:S30-S43.
26. Gehling M, Tryba M. Risks and side-effects of intrathecal morphine combined with spinal anaesthesia: a meta-analysis. *Anaesthesia* 2009;64(6):643-51.
27. Fleron MH, Weiskopf RB, Mouren S, Eyraud D, Godet G, Riou B, et al. A comparison of intrathecal opioid and intravenous analgesia for the incidence of cardiovascular, respiratory, and renal complications after abdominal aortic surgery. *Anesth Analg*. 2003;97(1):2-12.
28. Ko S, Goldstein DH, VanDerKerkhof EG. Definitions of "respiration depression" with intrathecal morphine postoperative analgesia: a review of literature". *Can J Anaesth*. 2003;50(7):679-88.
29. Etches RC, Sandler AN, Daley MD. Respiratory depression and spinal opioids. *Can J Anaesth*. 1989;36(2):165-85.
30. American Society of Anesthesiologist. Practice Guidelines for the Prevention, Detection and Management of Respiratory Depression Associated with Neuraxial Opioid Administration. *Anesthesiology* 2009:110.
31. Szarvas S, Harmon D, Murphy D. Neuraxial opioid-induced pruritus: a review. *J Clin Anesth*. 2003;15:234-9.

32. Jacobson L, Chabal C, Brody MC. A dose-response study of intrathecal morphine efficacy, duration, optimal dose, and side effects. . *Anesth Analg*. 1988;67:1082-8.
33. Borgeat A, Ekatodramis G, Schenker CA. Postoperative nausea and vomiting in regional anesthesia: a review. . *Anesthesiology*. 2003;98:530-47.
34. Ruslan IR, Murthi B, Judin M. Pengaruh Penambahan Morfin 0,1 mg, 0,15 mg dan 0,2 mg Intratekal Terhadap Analgesia dan Mual-Muntah Pasca Operasi Pada Anestesi Spinal: Program Studi Magister Kedokteran Keluarga Program Pascasarjana Universitas Sebelas Maret Surakarta; 2012 [cited 2015 18 Juli 2015]. Available from: <http://pasca.uns.ac.id/?p=2334>.
35. Mackintosh C. Assessment and management of patients with post-operative pain. *Nursing Standard*. 2007;22(5):49-55.
36. McCaffery M, Beebe A. *Pain: clinical manual for nursing practice*. St. Louis: C.V. Mosby; 1989. 353p p.
37. Carr EC, Nicky TV, Wilson BJ. Patient experiences of anxiety, depression and acute pain afer surgery: a longitudinal perspective. *Int J Nurs Stud*. 2005;42(5):521-30.
38. Wuhrman E, Cooney MF. *Acute Pain: Assessment and Treatment*: WebMD LLC; 2013 [cited 2014 4.20]. Available from: http://www.medscape.com/viewarticle/735034_3.
39. Tennant F. The physiologic effects of pain on the endocrine system. *Pain Ther*. 2013;2:75-86.
40. Onur E, Ercal T, Karslioglu I. Prolactin and cortisol level during spontaneous and oxytocin induces labour and the effect of meperidine. *Arch Gynecol Obstet*. 1989;244(4):227-32.
41. Benfield RD, Newton ER, Tanner CJ, Heitkemper MM. Cortisol as a biomarker of stress in term human labour: physiological and methodological issues. *Biol Res Nurs*. 2014;16(1):64-71.
42. American Society of Anesthesiologist Task Force on Chronic Pain Management. *Practice Guidelines for Chronic Pain Management*. *Anesthesiology* 2010:112.
43. Nikolajsen L, Sorensen H.C., Jensen T.S., Kehlet H. Chronic Pain following caesarean section. *Acta Anaesthesiol Scand*. 2004;48:111-6.
44. Eisenach J.C., Pan P.H., Smiley R., Lavandhomme P., Landau R., Houle T.T. Severity of acute pain after childbirth, but not type of delivery, predicts persistent pain and post-partum depression. *Pain*. 2008;140:87-94.
45. Brook P, Connel J, Pickering T. *Oxford Handbook of Pain Management*2011.
46. American Society of Anesthesiologists. *ASA physical status classification system* Washington DC: American Society of Anesthesiologists; 2014 [cited 2016 23-3-2016]. Available from: <https://www.asahq.org/resources/clinical-information/asa-physical-status-classification-system>.
47. Sastroasmoro S, Ismael S. *Dasar-dasar Metodologi Penelitian Klinis*. 4 ed. Jakarta: CV. sagung Seto; 2011. 358-9 p.

48. Lee HJ, Park JH, Kil HK, Choi SH, Noh SH, Koo BN. Efficacy of intrathecal morphine combined with intravenous analgesia versus thoracic epidural analgesia after gastrectomy. *Yonsei Med J.* 2014;55(4):1106-14.
49. Vadivelu N, Mitra S, Narayan D. Recent advances in postoperative pain management. *Yale Journal of Biology and Medicine* 2010;83:11-25.
50. Wong JY, Carvalho B, Riley ET. Intrathecal morphine 100 and 200 mcg for post cesarean delivery analgesia: a trade off between analgesic efficacy and side effects. *International Journal of Obstetric Anesthesia.* 2013;22(2):36-41.
51. DeSousa KA, Chandran R. Intrathecal morphine for postoperative analgesia: current trends. *World J Anesthesiol.* 2014;27(3):191-202.
52. Irawan H, Wahyudi, Hisbullah. Perbandingan kadar kortisol dan efek analgesia pascabedah anestesi spinal kombinasi bupivakain hiperbarik 0.5% 8 mg dan klonidin 30 mcg dengan bupivakain hiperbarik 0,5% 8 mg dan morfin 0,1 mg pada pasien yang menjalani prosedur seksio sesarea. *Anesthesia & Critical Care.* 2014;22(2):111-8.
53. Shafer SL, Varvel JR, Aziz N, Scott JC. Pharmacokinetics of fentanyl administered by computer-controlled infusion pump. *Anesthesiology.* 1990;73:1091-102.
54. Choi SH, Koo BN, Nam SH, Lee SJ, Kim KJ, Kil HK. Comparison of remifentanyl and fentanyl for postoperative pain control after abdominal hysterectomy. *Yonsei Med J.* 2008;49(204-10).
55. Burton D, Nicholson G, Hall G. Endocrine and metabolic response to surgery. *Continuing education in anesthesia, critical care & pain.* 2004;4(5):144-7.
56. Presseau EV, Martel MO, Mathieu R, Caron E, Albouy G, Marin MF, et al. Acute stress contributes to individual differences in pain and pain-related brain activity in healthy and chronic pain patients. *The Journal of Neuroscience.* 2013;33(16):6826-33.